

Stgr Wk. Cat. Lp. Eur. 296 no. 199 sp. 1997-8 (1871); Snln. Vlind. Nederl. Micr. 429, 669-71 no. 45 sp. 1-2 Pf. 8 · 268-9 (1882); Stgr-Rbl. Cat. Lp. Pal. 2 155 no. 315 sp. 2873-4 (1901); Busck Bull. US. Nat. Mus. 52 500-1 sp. 5596-605 (1902); Pr. US. Nat. Mus. 25 772, 807-15 Pf. 29 · 15 (1903); 30 725-6 tf. 726 · 4 (1906); Can. Ent. 38 212 (1906); Ent. News 19 316-7 (1908); Splr. Schm. Eur. 2 356-7 no. 16 sp. 1-2 (1910).

Type 2: *Evagora apicitripunctella* Clms. (Clms. 1860).

§*EVAGORA* Clms. Pr. Ac. Nat. Sc. Phil. 12 165-6 (1860); Clms-Stn. Tin. N-Am. x, 119-21 (1872); Wlsm. Tr. Am. Ent. Soc. 10 182 (1882); Pr. Z. Soc. Lond. 1897 63-6 sp. 21-6 (1897); Busck Pr. US. Nat. Mus. 25 807-9, 934 (1903); Pr. Ent. Soc. Wash. 5 200 (1903).

Type 3: *Gelechia* (*Teleia*?) *dorsivittella* Z. (= *vagatioella* Chmb.; Chmb. 1873).

EIDOTHEA Chmb. †*EIDOTHOA* Chmb. Can. Ent. 5 186-7 (1873). ‡*EIDOTHEA* Chmb. Can. Ent. 5 229-30 (1873). *EIDOTHOA* Busck Pr. US. Nat. Mus. 25 807-8, 812-3 (1903).

Type 4: *Anacamptis robiniella* Fitch (= *fuscopallidella* Chmb.; Chmb. 1873).

SINOE Chmb. Can. Ent. 5 229-231 (1873); Busck Pr. US. Nat. Mus. 25 807-8, 812-13 (1903).

Type 5: *Phalaena Tinea leucatella* Cl., L.

**RECURVARIA* (Hw.) HS. SB. Schm. Eur. 5 11, 44 no. 71 Pf. 14 · d (1853), 205-6 no. 71 sp. 573-4 (1854). *APHANAULA* Meyr. HB. Br. Lp. 569, 579-80 no. 7 sp. 1-2 (1895); Busck Pr. US. Nat. Mus. 25 807-8 (1903).

[**RECURVARIA* (Hw.) Meyr. HB. Br. Lp. 569, 606-7 no. 17 sp. 1 (1895) = *ACOMPSIA* Hb. (Type *cinerella* Cl.; Dp. 1838) *vide* Wlsm. Pr. Z. Soc. Lond. 1897 82 no. 18 (1897).]

1. *Recurvaria rhicnota*, sp. n. (Tab. II. fig. 4.)

Antennae pale cinereous, annulated with dark brown. *Palpi* with the median joint somewhat densely clothed, not hirsute, terminal joint as long as the median, acuminate; whitish cinereous, with two broad bands of dark brown on each joint, the anteterminal band on the median joint narrower than the other three. *Head* whitish cinereous, sprinkled with dark brown. *Thorax* dark brown, sprinkled with pale cinereous. *Forewings* olivaceous brownish, with a slight ochreous tinge, profusely speckled with pale cinereous, especially on their basal half; with numerous dark brown spots and blotches, of which the more conspicuous are two costal on either side of the middle, one discal a little beyond the middle, one dorsal before the tornus, and three or four smaller marginal, around the apex; cilia brownish ochreous, with a shade of fuscous speckling running through their middle. *Exp. al.* 15 mm. *Hindwings* narrower than the forewings, slightly bisinuate below the apex; bronzy brownish grey; cilia (1½) brownish grey. *Abdomen* greyish brown. *Legs*, hind tibiae with brownish ochreous hairs above; the tarsi greyish brown, with four pale annulations.

Type ♀ (65784) Mus. Wlsm. (*Godm-Salv. Coll.*) BM.

Hab. GUATEMALA: Las Mercedes, 3000 ft. (*G. C. Champion*). Unique.

Allied to 5598 *colubrinae* Busck.

2. *Recurvaria sartor*, sp. n. (Tab. II. fig. 7.)

Antennae black, with indistinct whitish annulations. *Palpi* recurved, median joint slightly roughened beneath, terminal nearly as long as median, slender; white, with two black annulations on the terminal,