

and a few black scales on the outer side of the median. *Head* white. *Thorax* pale fawn-brownish. *Forewings* fawn-brownish, with a lilac tinge (possibly due to abrasion), the costa is slightly shaded along the basal half; there is a patch of black raised scales on the dorsum at one-fourth, partially connected by blackish scaling along the dorsum, with a second smaller but otherwise similar spot at about two-thirds; cilia fawn-brownish about the apex, with a slight speckled appearance, becoming pale brownish grey on the tornus and at the dorsum. *Exp. al.* 11 mm. *Hindwings* pale shining silvery grey; cilia pale brownish grey. *Abdomen* dark brownish grey. *Legs* whitish, with four black tarsal spots above.

Type ♂ (65792) Mus. Wlsm. (*Godm-Salv. Coll.*) BM.

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Unique.

A distinct species.

3. *Recurvaria thysanota*, sp. n. (Tab. II. fig. 5.)

Antennae somewhat stout; cinereous, with faint darker annulations. *Palpi* chalky-white, the median joint somewhat expanded on either side at its apex, shaded with brownish fuscous externally at its base; terminal joint about as long as the median, porrect, with two fuscous annulations. *Head* and *Thorax* chalky-white, the tegulae smeared with brownish fuscous. *Forewings* chalky-white, somewhat shaded and sprinkled with brownish fuscous; a dark brownish fuscous line on the base along the fold and another from the middle of the wing to above the apex are blended by brownish shading on the basal half of the cell; a small fuscous dot lies a little above the outer extremity of the fold, and there is another at the extreme apex; apical and subapical cilia whitish, sprinkled with brownish fuscous; tornal cilia unsprinkled, inclining to brownish ochreous. *Exp. al.* 10 mm. *Hindwings* nearly as broad as the forewings, oblique below the produced apex; ♂ with a conspicuous yellowish ochreous limbal hair-pencil; shining, pale steely grey; cilia brownish cinereous, with a slight ochreous tinge. *Abdomen* shining, steel-whitish. *Legs* whitish cinereous.

Type ♂ (65786) Mus. Wlsm. (*Godm-Salv. Coll.*) BM.

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Unique.

Most nearly allied to *juniperella* Krft. [*Jr. N-Y. Ent. Soc.* 11 153, 157-8 Pf. 9·3, 17 (1903)], but separable by the limbal hair-pencil, and the slightly different markings.

4. *Recurvaria flagellifer*, sp. n.

Antennae pale brownish fuscous. *Palpi* white, the median joint black nearly to its apex on the outer side, but not within nor above; terminal joint with two blackish annulations. *Head* and *Thorax* dirty whitish, the latter sprinkled with brownish fuscous. *Forewings* dirty whitish, profusely sprinkled with brownish fuscous, with numerous small spots of pale, slightly raised scales; the fuscous clouding tends to assume the form of oblique shades descending outward from the costa, but these are so confused and ill-defined that they do not constitute true bands or fasciae, nor do they reach the dorsum; the first of these is near the base and can be traced to the fold and a little beyond it; the second is a little before the middle, rather less oblique, but in some specimens more strongly marked than the first; the third consists mainly of a larger dark fuscous cloud before the commencement of the costal cilia, and opposite to this is a smaller patch on the dorsum, the terminal portion of the wing being also more or less suffused; cilia of the same colour as the wing-surface, with the same speckled appearance. *Exp. al.* 11-13 mm. *Hindwings* shining, pale greyish; cilia shining, brownish ochreous; ♂ with strong bright ochreous flexal hair-pencil. *Abdomen* brownish fuscous. *Legs* dirty whitish, with fuscous tarsal shade-bands.

Type ♂ (65777); ♀ (65778) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PTT.* (65779-80) US. Nat. Mus.]

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Seven specimens.