




J. H. Durrant, del.

12. *Arogalea cristifasciella* Chmb. ♀.

Nearest to *Telphusa* Chmb., from which it differs in having HW: 6-7 separate, and from *Leptogeneia* Meyr. [Pr. Linn. Soc. NSW. 29 257, 412-3 no. 47 (1904)] in the lack of the rough projecting tuft on the lower side of the palpi, as also in having FW: 2 remote from 3. Although not nearly related, *Arogalea* is liable to be confused in tabulations with *Aristotelia* Hb., owing to the separation of HW: 6-7. *Arogalea* differs in these veins not being truly parallel, in HW: 5 being bent towards 4 at base, and in the presence of raised scales in the forewings.

1. *Arogalea archaea*, sp. n. (Tab. II. fig. 10.)

Antennae white, delicately annulate with brown. *Palpi* white, the median joint slightly thickened, with a broad olive-brown band near the base, a narrower one near its apex; terminal joint with a narrow band near its base, a wide one at its apex. *Head* and face snow-white. *Thorax* white, dusted with olive-brown. *Forewings* white, profusely dusted with olive-brown, with many brownish fuscous spots and mottlings; a small one at the base of the costa is followed by another a little beyond it, a third lying below the base of the fold; a dorsal streak at about one-fifth crosses the fold upward and terminates on the cell; above and beyond this are two more spots, one on the costa, one below and beyond it; these are followed by a broad brownish fuscous costal blotch, beyond the middle, which merges at its lower end in an olive-brown median shade, and opposite to it is a scarcely smaller dorsal blotch, the apical portion of the wing being mottled throughout with brownish fuscous, mixed with olive-brown; cilia white, with lines of olive-brown dusting running through them, toral cilia pale brownish cinereous. *Exp. al.* 13 mm. *Hindwings* (1) shining, pale grey; cilia brownish cinereous. *Abdomen* brownish cinereous. *Legs* pale brownish cinereous, the tarsal joints distinctly banded with fuscous.

Type ♂ (65794) Mus. Wlsm. (Godm-Salv. Coll.) BM.

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII. (H. H. Smith). Unique.

2. *Arogalea senecta*, sp. n.

Antennae stout, greyish fuscous. *Palpi* white, with a broad blackish band at the base and a narrower one before the apex of the median joint; with two blackish bands around the terminal joint, one before, the other beyond the middle. *Head* and *Thorax* greyish white, sprinkled with dark fuscous. *Forewings* narrow, elongate, greyish white, with dark fuscous sprinkling throughout and black scale-patches, some of which are distinctly raised; a small raised black spot on the costa near the base, a larger black spot in and above the fold at about one-fourth, a distinctly raised black patch on and above the dorsum beneath it, accompanied by a few chestnut-brown scales, also raised; beyond this series of spots, which are more or less