

connected, a black costal spot occurs before the middle, with a curved black streak a little below and beyond it at the upper edge of the cell; this is succeeded by an inwardly oblique blackish cloud coming from the costa beyond the middle and almost connected with an opposite black dorsal spot, the latter narrowly produced outward and diffused nearly to the tornus; the apical portion of the wing thickly sprinkled with dark fuscous, a streak of the same running through the basal half of the cilia at the apex; cilia whitish grey, assuming an ochreous tint along the dorsum. *Exp. al.* 12 mm. *Hindwings* pale, shining, bluish grey; cilia ochreous grey. *Abdomen* shining, pale brownish ochreous. *Legs*: anterior pair barred with black and white; posterior legs whitish, the tarsal joints with blackish annulations.

*Type* ♂ (65797) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (65798) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., IX. (*H. H. Smith*). Two specimens.

### 3. *Arogalea albilingua*, sp. n.

*Antennae* white, with faint darker annulations. *Palpi* with strong sooty black patch at the base of the median joint, a small brown spot before the apex; with a black annulus near the base of the terminal joint and a wider one near its apex. *Haustellum* scaled with white at its base, showing somewhat conspicuously between the dark eyes and bases of the palpi. *Maxillary Palpi* short, whitish. *Head* shining, silvery white. *Thorax* white, faintly dusted with brownish fuscous. *Forewings* shining white, sprinkled with brownish ochreous and brownish fuscous scales; with five conspicuous black marginal spots, one of these at the base of the costa is reduplicated, the white ground showing between its extremities; a second costal spot, before the middle, is followed by a third beyond the middle, opposite to which and a little beyond it is a black dorsal spot of raised scales at the end of the fold; beyond this again is a speck of black scales on the termen a little above the tornus; another conspicuous black spot, with raised white scales intermixed, lies across the middle of the fold, a smaller similar spot on the cell about the middle of the wing; cilia white, with some greyish fuscous dusting along their base, becoming shining ochreous on the dorsum; underside fuscous. *Exp. al.* 10–10.5 mm. *Hindwings* pale steely grey; cilia shining, ochreous; underside pale rosy grey. *Abdomen* shining, pale greyish ochreous. *Legs* shining, whitish ochreous, the hind tarsi faintly spotted.


*Type* ♂ (65795) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (65796) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Two specimens.

### ZELOSYNE, gen. n.

(ζηλοσύνη = emulation.)

*Type*: *Zelosyne poecilosoma* Wlsm. (Tab. II. fig. 11.)



J. H. Durrant, del.

13. *Zelosyne poecilosoma* Wlsm. ♂.