





65806.



J. H. Durrant, del.

15. *Eudactylota barberella* Busck. ♀.

curve from 6 to flexus: *neuration* 12 veins; 7-8 stalked, to costa; rest separate, 2 more remote from 3 than 3 is from 4. *Hindwings* 1, bilobed in both ♂ and ♀, but slightly differing in shape; costa nearly straight to apex which is pointed; termen excised between 6 and 5 and forming two lobes, the apical long and pointed, the lower much shorter, broad, its somewhat blunt apex being slightly above 5; dorsum somewhat sinuate between 1<sup>a</sup> and 2: *neuration* 8 veins; 6-7 stalked; 3-4 separate; 5 cubital; discoidal subobsolete on fold above 5; 12 connected to radius by 11. *Abdomen* moderate. *Legs*: hind tibiae hairy.

Allied to *Neodactylota* Busck, but HW:5 is separate from 4; the hindwings are bilobed in both sexes, and the ♂ lacks the pouch containing a hair-pencil.

### 1. *Eudactylota barberella* Busck. (Tab. II. fig. 13.)

*Neodactylota barberella* Busck [Bull. US. Nat. Mus. 52 504 sp. 5644 (1902)<sup>1</sup> LN.]: Pr. US. Nat. Mus. 25 835, 836-7 Pf. 30 · 23 (1903)<sup>2</sup>; Krft. Smith's Check List Lp. Bor-Am. 110 sp. 6190 (1903)<sup>3</sup>.

*Antennae* cinereous, annulate with black, the annulations connected by slender black lines above. *Palpi* smooth, recurved to above the vertex, terminal joint longer than the median; cinereous, dusted with brownish fuscous scales. *Head* and *Thorax* greyish brown. *Forewings* tawny chestnut-brown, dusted with pale cinereous and fuscous scales, especially on a transverse fascia from costa to tornus at two-thirds of the wing-length; the lower half of this fascia is margined on its inner side by a triangular dark fuscous shade, and the costa is narrowly shaded with fuscous from the base to the upper extremity of the fascia; cilia tawny, dusted with fuscous atoms. *Exp. al.* 11-12 mm. *Hindwings* cinereous; cilia with an aeneous gloss. *Abdomen* shining, greyish cinereous; anal tuft inclining to ochreous. *Hind Legs* fuscous, speckled with pale cinereous scales, with two pale bands around the tibiae, and three around the tarsal joints.

*Type* ♂ (6367) US. Nat. Mus. [Nt. ♀ (65801 *Amula*) Mus. Wlsm. (*Godm-Salv. Coll.*) BM.]

*Hab.* AMERICA, N-C. **United States**<sup>1-3</sup>—ARIZONA: Williams, 27. V. 1901 (*H. S. Barber*)<sup>2</sup> [= "Col."<sup>1</sup>]. **Central America**—MEXICO: GUERRERO: *Amula*, 6000 ft., VIII-IX. (*H. H. Smith*). Six specimens.