

5. *Telphusa niveisignella* Z.

Psoricoptera niveisignella Z. Hor. Soc. Ent. Ross. 13 333-5 Pf. 4·106 (1877)¹.

Hab. AMERICA, C-S. Central America¹—PANAMA¹: CHIRIQUI¹ (*Ribbe*). South America—COLOMBIA: CUNDINAMARCA: Bogota (*Baron von Nolcken*)—PERU: CUZCO: Cuzco Mts. (*Garlepp*, Stgr. 1895).

The specimen from Bogota was determined by Zeller, and agrees with *chionostigma* in having HW:6-7 stalked, and HW:3-4 remote, as also in the clothing of the palpi. The Cuzco specimen (73496) is in much finer condition, and the terminal joint of the palpi is consequently more evenly and strongly clothed.

6. *Telphusa glandiferella* Z.

=*sella* Chmb.³; =†*glandifuella* Chmb.; =†*glandifera* Chmb.

Gelechia (*Anacampsis*) *glandiferella* Z. Verh. ZB. Ges. Wien 23:1873 275-6 Pf. 4·25 (1873)¹.

Gelechia sella Chmb. Can. Ent. 6 238 (1874)². *Gelechia glandifuella* Chmb. Can. Ent. 9 14

(1877)³. *Gelechia glandifera* Chmb. Can. Ent. 9 24 (1877)⁴. *Gelechia glandiferella* Chmb.

Bull. US. GG. Surv. 4 144 (1878)⁵; Hgn. Pap. 4 99 (1884)⁶. *Gelechia sella* Frey Pap. 4 99

(1884)⁷. *Gelechia glandiferella* Riley, Smith's List Lp. Bor-Am. 101 sp. 5382 (1891)⁸.

Xenolechia glandiferella Wlsm. Pr. Z. Soc. Lond. 1897 72 sp. 38 (1897)⁹. *Telphusa glandi-*

ferella Busck Bull. US. Nat. Mus. 52 497 sp. 5564 (1902)¹⁰; Pr. US. Nat. Mus. 25 784,

788-9 (1903)¹¹; Krt. Smith's Check List Lp. Bor-Am. 109 sp. 6105 (1903)¹².

Hab. AMERICA, N-C-I. United States¹⁻¹²—KENTUCKY⁷—TEXAS^{1-6, 9-12}: VII-VIII¹: BOSQUE CO.: 11-20. V, 22. VII, 10-25. X. (*G. W. Belfrage*)—CALIFORNIA: MENDOCINO: Head of Noyo, 8-11. VI. 1871 (*Wlsm.*): SHASTA: Hatchet Creek, 14-17. VII. 1871 (*Wlsm.*); Pit River, 21-26. VII. 1871 (*Wlsm.*): SISKIYOU: Mt. Shasta, 2.VIII-1.IX. 1871 (*Wlsm.*). Central America—MEXICO: TABASCO: Teapa, II. (*H. H. Smith*). West Indies⁹—GRENADA⁹: Mt. Gay Estate, 3000 ft., VIII. (*H. H. Smith*)⁹.

PSORICOPTERA Stn.

Type: *Gelechia* (*Chelaria*) *gibbosella* Z. (Stn. 1854).

PSORICOPTERA Stn. Ins. Br. Lp-Tin. (†*PHORICOPTERA* 76), 100-1 no. 7 sp. 1 Pf. 4·4^c (1854); Hnmn. Schm. Deutsch. Tin. 188, 192 no. 417 sp. 270 (1871), Tbl. 5, 31 no. 417 (1876); Stgr-Wk. Cat. Lp. Eur. 289 no. 193 sp. 1788 (1871); Snln. Vlind. Nederl. Micr. 428, 609-11 no. 41 sp. 1 Pf. 8·259, 272 (1882); Meyr. HB. Br. Lp. 569, 604-5 no. 14 sp. 1 (1895); Stgr-Rbl. Cat. Lp. Pal. 2 140 no. 299 sp. 2507-8 (1901); Splr. Schm. Eur. 2 359 no. 23 Pf. 87·60 (1910).

1. *Psoricoptera trigonota*, sp. n.

Antennae slightly biserrate toward the apex; umber-brown. *Palpi* strongly recurved, the terminal joint longer than the median, closely clothed above to the apex; cinereous, mottled with umber-brown. *Head* dark umber-brown; face shining, fawn-ochreous. *Thorax* dark brown, mottled with shining fawn-ochreous. *Forewings* shining, fawn-ochreous, mottled with darker and paler shades of umber-brown, with some fuscous scaling at the base; on the commencement of the cell is a conspicuous flap of scales;