

## 2. *Gelechia phalacra*, sp. n.

*Antennae* blackish, not annulate. *Palpi* cream-white, the median joint with an even brush beneath; terminal joint tipped with fuscous. *Head* cream-coloured, with long white scales meeting over the vertex. *Thorax* cream-coloured, tegulae dark fuscous. *Forewings* with straight margins and lanceolate apex; dark fuscous, a slender streak of pale cream-colour from the base along the margin to the flexus; a small costal spot, at the commencement of the cilia, preceded by a larger dorsal spot, both pale cream-colour; a minute spot of the same on the cell, about the middle of the wing, is preceded by another at the apex, and others, along the termen, are followed by black scales; a similar spot lies on the fold at about its middle; cilia dark fuscous. *Exp. al.* 9 mm. *Hindwings* pale greyish cinereous; cilia slightly tinged with fuscous. *Abdomen* greyish cinereous, tinged with fuscous. *Legs* fuscous, with pale cream-coloured annulations.

*Type* ♂ (35653) Mus. Wlsm. BM. [PT. (35654) US. Nat. Mus.]

*Hab.* MEXICO: SONORA: 1883 (*H. K. Morrison*). Two specimens.

Most nearly allied to *donatella* Wkr., but the pale spots are smaller and the terminal joint of the palpi is slightly longer and more shaded.

## 3. *Gelechia lacticaput*, sp. n.

*Antennae* dark brown, cream-colour at the base to one-fifth of their length, with some creamy speckling near their apex. *Palpi* cream-colour, the median joint somewhat thickened, the terminal joint with two slight brown annulations. *Head* and *Thorax* cream-colour, the latter with a dark brown patch posteriorly. *Forewings* dark brown, with a very narrow snow-white basal patch extending across the wing, an oblique snow-white costal patch at one-fourth reaching to the fold and projecting outward on the cell, a minute white oblique costal streak at the commencement of the costal cilia, and a large snow-white semicircular tornal patch extending through the terminal cilia and sparsely sprinkled with brownish scales; cilia at the apex dark brown. *Exp. al.* 15 mm. *Hindwings* pale purplish grey; cilia with a slight ochreous tinge. *Abdomen* grey. *Legs* creamy; tarsi banded and speckled externally with dark brown.

*Type* ♂ (65682) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PT. (65683) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Two specimens.

A conspicuous and distinct species.

## 4. *Gelechia cerussata*, sp. n. (Tab. II. fig. 18.)

*Antennae* fuscous, annulate with pale cream-ochreous. *Palpi* pale cream-ochreous; the median joint moderately thickened, mottled with fuscous on the outer side and beneath; terminal joint with two fuscous annulations. *Head* and *Thorax* pale cream-ochreous. *Forewings* obtusely pointed; dark purplish fuscous, with a very narrow, pale cream-ochreous, straight, transverse band at the extreme base, and an elongate-ovate white patch, sparsely dusted with fuscous, occupying the tornus and termen, with the cilia, to below the apex, near which it encloses a dark fuscous dot; the cilia at the extreme apex and above it smoky fuscous; a blackish dot occurs on the disc, above the middle, at scarcely one-half from the base. *Exp. al.* 14 mm. *Hindwings* grey; cilia pale brownish grey. *Abdomen* (missing). *Legs* pale cream-ochreous; the hind tibiae twice banded obliquely with smoky grey externally; hind tarsi sprinkled externally with fuscous.

*Type* ♂ (5977 Wlsm. Det. 1908) US. Nat. Mus.

*Hab.* MEXICO: VERA CRUZ: Cordova, V. 1908 (*F. Knab*). Unique.

## 5. *Gelechia infracta*, sp. n.

*Antennae* fuscous. *Palpi* dark fuscous, the median joint with two, the terminal joint with three paler