

subochreous bands, the last including the apex. *Head* and *Thorax* dark fuscous; a few ochreous scales above the eyes. *Forewings* tawny fuscous, with a dark fuscous median shade running through them, from near the base to the apex, interrupted before the middle by a large white, outwardly oblique, costal patch which reaches to the fold expanding outward on the cell; this is clearly defined, its origin being at one-fourth from the base; there is a faintly indicated spot of pale scales at the end of the cell and another above and beyond it on the costa; the dorsum is narrowly white from base to flexus; cilia greyish fuscous, sprinkled with whitish ochreous. *Exp. al.* 17 mm. *Hindwings* shining, pale grey; cilia inclining to ochreous. *Abdomen* leaden grey. *Legs* whitish ochreous, the tarsi and tibiae widely banded with fuscous.

*Type* ♂ (65684) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (65685) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Two specimens.

### 6. *Gelechia cacoderma*, sp. n.

*Antennae* tawny fuscous, faintly annulate with pale ochreous. *Palpi* tawny fuscous, speckled with pale ochreous. *Head* whitish ochreous, speckled with tawny fuscous. *Thorax* tawny fuscous, speckled with pale ochreous. *Forewings* tawny fuscous, intermingled with pale brownish ochreous and blotched with brownish fuscous, the usual markings being scarcely distinguishable in the somewhat rough and mixed scaling; the darker blotching apparently coincides with a medio-plical spot, a discal spot beyond it, and another at the end of the cell, and forms a costal spot at one-fourth, another at three-fifths, and an apical shade preceded by an ill-defined transverse pale band; cilia pale brownish ochreous, speckled with tawny fuscous along their basal half. *Exp. al.* 16-21 mm. *Hindwings* broader than the forewings, scarcely sinuate below the apex; bronzy brownish, with paler greyish brown cilia. *Abdomen* brownish ochreous towards the base, shading to greyish brown posteriorly. *Legs* greyish brown, hind tibiae and tarsi banded with pale ochreous.

*Type* ♂ (65376); ♀ (65377) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PTT.* (65381-2, 65385) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Twelve specimens.

Closely allied to 5755 *bicostimaculella* Chmb. in colouration, but a much larger species with the paler markings less distinct.

### 7. *Gelechia veneranda*, sp. n.

*Antennae* fawn-brownish, delicately barred with blackish above. *Palpi* brownish fawn, streaked with blackish, the terminal joint nearly black throughout; the median very hirsute beneath, with long diverging scales. *Head* stone-whitish. *Thorax* fawn-grey. *Forewings* mottled with much mixed colours, of which the ground-colour may be taken as pale stone-colour; on this is a profuse sprinkling and partial suffusion of fawn-brownish, with sundry spots and streaks almost black, or at least very dark fuscous; of these two are costal, one before and one beyond the middle, and two are discal, parallel to the costal, but a little further removed from the base; the terminal and apical portion of the wing is almost entirely suffused with this dark brownish fuscous shading, and the fold is marked by a narrow line of the same throughout; all the different shades are much blended together and ill-defined, a slight rufous tinge prevailing above the fold; cilia brownish ochreous, the outer half paler. *Exp. al.* 16-17 mm. *Hindwings* brownish grey, profusely irrorated with brownish scales beyond the basal third; cilia pale brownish grey. *Abdomen* brownish cinereous. *Legs* brownish fuscous, with pale cinereous spots on the tarsal joints.

*Type* ♂ (35655); ♀ (35656) Mus. Wlsm. BM. [*PTT.* (35659, 35661-2) US. Nat. Mus.]

*Hab.* MEXICO: SONORA: 1883 (*H. K. Morrison*). Eight specimens.