

pp. ix, 218, 224 (1872)². *Gelechia vagella* Wkr. Cat. Lp. BM. 29 596 sp. 237 (1864)³. *Depressaria fuscoochrella* Chmb. Can. Ent. 4 106, 129, 147-8 (1872)⁴. *Gelechia (Lita) liturosella* Z. Verh. ZB. Ges. Wien 23:1873 265 (1873)⁵. *Gelechia fuscoochrella* Chmb. Bull. US. GG. Surv. 4 143 (1878)⁶. *Gelechia liturosella* Chmb. Bull. US. GG. Surv. 4 144 (1878)⁷. *Gelechia mediofuscella* Chmb. Bull. US. GG. Surv. 4 144 (1878)⁸. *Gelechia (Lita) vagella* Wlsm. Tr. Am. Ent. Soc. 10 178 (1882)⁹. *Gelechia fuscoochrella* Hgn-Frey Pap. 4 98 (1884)¹⁰. *Gelechia liturosella* Hgn. Pap. 4 99 (1884)¹¹. *Gelechia mediofuscella* Riley, Smith's List Lp. Bor-Am. 101 sp. 5407 (1891)¹². *Gelechia vagella* Riley, Smith's List Lp. Bor-Am. 102 sp. 5506 (1891)¹³. *Gelechia mediofuscella* Busck Bull. US. Nat. Mus. 52 514 sp. 5764 (1902)¹⁴; Pr. US. Nat. Mus. 25 854, 885, 935 (1903)¹⁵; 27 759 (1904)¹⁶; Pr. Ent. Soc. Wash. 5 211 (1903)¹⁷; Krft. Smith's Check List Lp. Bor-Am. 112 sp. 6268 (+9268) (1903)¹⁸.

Hab. AMERICA, N-C. Canada³—NOVA SCOTIA³. United States^{1-2, 4-18}—ATLANTIC STATES¹⁴—MASSACHUSETTS⁹—DISTRICT OF COLOMBIA¹⁵—NORTH CAROLINA: 1883 (*H. K. Morrison*)—KENTUCKY^{4, 10, 14}—MISSOURI: ST. LOUIS: Kirkwood, 1883-4 (*M. Murtfeldt*)—TEXAS^{5, 11, 14}—WASHINGTON¹⁶. Central America—MEXICO: SONORA: 1883 (*H. K. Morrison*): VERA CRUZ: Cordova, V. 1908 (*F. Knab*): GUERRERO: Amula, 6000 ft., IX. (*H. H. Smith*)—PANAMA: CHIRIQUI: Volcan de Chiriqui, 2000-3000 ft., 1881 (*G. C. Champion*).

15. *Gelechia repentina*, sp. n.

Antennae greyish fuscous. *Palpi* brownish ochreous, the median joint slightly roughened, with two slight bands of fuscous scales; terminal joint about as long as the median, also twice banded with fuscous. *Head* brownish ochreous. *Thorax* brownish ochreous, mottled with fuscous posteriorly. *Forewings* brownish ochreous, mottled with fuscous; a broad bluish fuscous band across their middle, occupying a space equal to half their length, its inner and outer margins both inclining to be concave and somewhat darker than the centre; a few reddish ochreous scales beyond the end of the cell are merged in a diffused fuscous shade extending to the apex; cilia pale ochreous, sprinkled with fuscous dots. *Exp. al.* 12-16 mm. *Hindwings* a little broader than the forewings; pale greyish brown, slightly transparent near their base; cilia pale brownish cinereous. *Abdomen* cinereous. *Legs* cinereous; the femora with one, the tibiae with two, and the tarsi with three brownish fuscous bands externally.

Type ♂ (65666); ♀ (65667) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PT. (65668-9) US. Nat. Mus.]

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Five specimens.

Closely allied to *mediofuscella* Clms., but the dark markings extend across the wing, and their inner edge seems to be less oblique and to commence slightly further from the base.

16. *Gelechia neptica*, sp. n.

Antennae yellowish brown, obscurely banded with fuscous. *Palpi* yellowish brown, median joint brush-like beneath, with two fuscous bands externally; terminal joint with a fuscous band before the apex. *Head* pale cinereous, speckled with fuscous. *Thorax* yellowish brown. *Forewings* yellowish brown, with some scattered dark fuscous scales; a dark fuscous spot at the base of the dorsum, and a small one in the middle of the fold, with another at the outer end of the fold; the first plical spot is followed by an