

oblique, elongate, very dark fuscous discal spot, a smaller costal spot preceding the cilia; the pale yellowish brown terminal cilia are much dusted with dark fuscous, which is thicker at and around the apex. *Exp. al.* 15 mm. *Hindwings* pale brownish grey; cilia brownish cinereous. *Abdomen* brownish cinereous above, tawny fuscous at the sides; pale ochreous beneath. *Legs* tawny fuscous, banded with very pale ochreous on the tibiae and around the tarsal joints.

*Type* ♂ (65671, *Omiteme*) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PTT. (65672, 65674) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Omiteme, 8000 ft., VII. (*H. H. Smith*); Amula, 6000 ft., VIII. (*H. H. Smith*). Four specimens.

### 17. *Gelechia bufo*, sp. n.

*Antennae* yellowish brown, with black annulations. *Palpi* hoary, the median joint barred with black externally, and sprinkled with black and yellowish brown beneath; the terminal joint with two blackish bands. *Head* hoary, sprinkled with yellowish brown scales. *Thorax* yellowish brown. *Forewings* elongate, narrow, lanceolate; yellowish brown, mottled with black; after a small black spot at the extreme base of the costa there is, at one-third, a triangular black costal spot, produced obliquely downward to the fold, and thence diffused in an overflow of blackish scales along the dorsum to a cloudy spot of the same before the tornus; there is also a smaller black spot on the middle of the costa, with another placed obliquely below it on the cell, and a third costal spot before the cilia, the space beyond the cell, as well as the terminal and apical area, being much clouded with thickly bestrewed blackish scales, some appearing also on the brownish ochreous cilia. *Exp. al.* 13-14 mm. *Hindwings* leaden grey; cilia brownish grey. *Abdomen* greyish fuscous. *Legs* hoary, blotched with brown externally.

*Type* ♂ (65662) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PTT. (65663, 65665) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Four specimens.

### 18. *Gelechia inaequalis*, sp. n.

*Antennae* brownish white, barred above with brownish fuscous. *Palpi* white, the terminal joint much dusted with brownish fuscous. *Head* white. *Thorax* brownish white, smeared with brownish fuscous anteriorly. *Forewings* brownish white, dusted with brownish fuscous, increasingly towards the apex; with two brownish fuscous incomplete fasciae—the first at one-third from the base, tending a little obliquely outward from the costa, its inner edge clearly defined, its outer side somewhat diffused toward a small discal spot opposite its middle; the second, much smaller and not oblique, descending from the costa at two-thirds, absorbing a small spot at the end of the cell and merging in the paler speckling of the surface below the cell; there is a small spot below the costa, half-way between the base and the first fascia, and one or two slight spots on the termen at the base of the brownish white, somewhat dusted, cilia. *Exp. al.* 10 mm. *Hindwings* grey; cilia pale greyish ochreous. *Abdomen* brownish grey. *Legs* whitish cinereous, with pale fuscous tarsal spots.

*Type* ♂ (35644) Mus. Wlsm. BM. [PTT. (35645-6) US. Nat. Mus.]

*Hab.* MEXICO: SONORA: 1883 (*H. K. Morrison*). Four specimens.

Somewhat resembling 5564 *Telphusa glandiferella* Z., but the conspicuous dark spot is costal, not dorsal.

### 19. *Gelechia concinna*, sp. n. (Tab. II. fig. 22.)

*Antennae* creamy white, delicately annulated with fuscous. *Palpi* creamy white, the median joint roughened beneath; the terminal joint as long as the median, sprinkled with fuscous, and banded before the apex. *Head* and *Thorax* creamy white, the latter somewhat shaded with greyish fuscous. *Forewings* creamy white, delicately shaded with brownish ochreous, especially below the fold and about the end of the cell; a minute fuscous spot at the base of the costa is followed by a triangular fuscous shade, of which the outer extremity scarcely reaches beyond the middle; a small fuscous spot at the