

grey, blending to bronzy brownish outward, especially toward the margins; cilia brownish cinereous. *Abdomen* brownish fuscous, with narrow pale segmental bands. *Legs* dark fuscous externally, pale cinereous on their inner sides.

*Type* ♂ (35637); ♀ (35638) Mus. Wlsm. BM. [PTT. (35645-6) US. Nat. Mus.]

*Hab.* MEXICO: SONORA: 1883 (*H. K. Morrison*). Seven specimens.

### 23. *Gelechia pleroma*, sp. n.

*Antennae* fuscous. *Palpi* with the median joint clothed with a rather short roughened brush beneath; fuscous, sprinkled with pale cinereous; the terminal joint scarcely shorter than the median. *Head* fuscous, mixed with pale cinereous; face pale brownish cinereous. *Thorax* purplish fuscous, touched with pale cinereous posteriorly. *Forewings* with the costa slightly arched at the base, straight beyond, apex slightly depressed, somewhat obtuse, termen oblique; purplish fuscous, with very faintly indicated pale brownish cinereous mottling, a small costal spot of the same at three-fourths from the base somewhat more distinct; cilia greyish fuscous. *Exp. al.* 16 mm. *Hindwings* bronzy brownish; cilia greyish brown. *Abdomen* fuscous above, with narrow pale transverse bars; anal tuft brownish cinereous. *Legs* pale brownish cinereous, banded externally with fuscous.

*Type* ♂ (65375, *Amula*) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PT. (5835 Wlsm. Det. 1908) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: *Amula*, 6000 ft., IX. (*H. H. Smith*): OAJACA: *Salina Cruz* (*W. Schaus*). Two specimens.

### 24. *Gelechia scotodes*, sp. n. (Tab. II. fig. 25.)

*Antennae* dark purplish fuscous. *Palpi* with a rough divided brush beneath, terminal joint not longer than the median; dark tawny fuscous, except along the upper surface of the median joint which is shining whitish ochreous. *Head* and *Thorax* smooth; dark tawny fuscous. *Forewings* with the apex depressed, obtusely rounded; dark tawny fuscous, obscurely sprinkled with pale cinereous in some lights; three spots darker than the ground-colour, two discal and one plical, the plical immediately below the first discal, accompanied by a few pale scales; a small whitish spot at the commencement of the costal cilia, without indication of a transverse fascia; cilia pale cinereous, much suffused with tawny fuscous. *Exp. al.* 14 mm. *Hindwings* bluish grey, overspread with brownish scales, more thinly along the cell and at the base; cilia brownish grey. *Abdomen* brownish fuscous. *Legs* whitish ochreous, suffused and banded with tawny fuscous.

*Type* ♂ (35628) Mus. Wlsm. BM.

*Hab.* MEXICO: SONORA: 1883 (*H. K. Morrison*). Unique.

### 25. *Gelechia hetaeria*, sp. n. (Tab. II. fig. 24.)

*Antennae* cinereous, fuscous above. *Palpi* cinereous, banded with black near the apex of the median joint; terminal joint suffused with black. *Forewings* rosy greyish fuscous, a black spot at the base of the costa and three elongate black spots, somewhat ill-defined, along the discal cell, with a small one beyond the end of the cell and a few black marginal spots around the apex; cilia rosy, slightly sprinkled with fuscous atoms. *Exp. al.* 15 mm. *Hindwings* pale leaden grey; cilia brownish grey, with a slight rosy tinge. *Abdomen* greyish fuscous. *Legs* cinereous, suffused and sprinkled with blackish scales.

*Type* ♂ (65688) Mus. Wlsm. (*Godm-Salv. Coll.*) BM.

*Hab.* MEXICO: VERA CRUZ: *Orizaba*, XII. 1887 (*F. D. Godman and O. Salvin*). Unique.