

towards the tornus, beyond which a sooty streak runs to the apex and through the cilia; the upper edge of the dark portion is slightly undulate and there is some admixture of ochreous scales upon its whole surface; cilia greyish fuscous, with a slight sprinkling of ochreous scales. *Exp. al.* 11 mm. *Hindwings* grey; cilia brownish grey. *Abdomen* brownish grey. *Legs* brownish fuscous, with pale ochreous annulations.

Type ♀ (35631) Mus. Wlsm. BM.

Hab. MEXICO: SONORA: 1883 (*H. K. Morrison*). Unique.

POLYHYMNO Chmb.

=COPOCERCIA Z.

Type 1: Polyhymno luteostrigella Chmb. (Chmb. 1874).

POLYHYMNO Chmb. Can. Ent. 6 246-9 (1874): 8 30-1 (1876): Bull. US. GG. Surv. 4 161 (1878): Jr. Cin. Soc. NH. 2 198, tf. 203·26 (1880); Riley, Smith's List Lp. Bor-Am. 104 sp. 5586-7 (1891); Wlsm. Tr. Ent. Soc. Lond. 1891 95-7 Pf. 4·32-3, 7·78 (1891): Pr. Z. Soc. Lond. 1891 525, 546 no. 49 sp. 90 (1891): 1897 77-9 no. 16 sp. 54-7 (1897); Busck Jr. NY. Ent. Soc. 8 235-6 Pf. 9·1 (1900): Bull. US. Nat. Mus. 52 504 sp. 5647-9 (1902): Pr. US. Nat. Mus. 25 773, 839-40 Pf. 31·26 (1903); Krtt. Smith's Check List Lp. Bor-Am. 110 sp. 6193-5 (1903); Meyr. Pr. Z. Soc. Lond. 1908 728 (1908): Ann. Transv. Mus. 2 14-16, Pf. 5·3-6 (1909).

Type 2: Copocercia crambinella Z. (Z. 1877).

COPOCERCIA Z. Hor. Soc. Ent. Ross. 13 374-7 Pf. 5·129^{a-b} (1877).

1. *Polyhymno subaequalis*, sp. n.

Antennae whitish, slightly tinged with bronzy brown. *Palpi* whitish. *Head* white. *Thorax* white, tinged with bronzy brown along the middle. *Forewings* shining bronzy brown, with a broad white median streak from the base, attenuated to a point beyond the end of the cell where it nearly coalesces with a small elongate white spot beneath its apex, which is succeeded by a white triangular patch before the termen; the lower angle of this is produced in a narrow marginal line along the base of the dorsal cilia, its upper angle also produced towards the apex, but shortly meeting, at an acute angle, two white costal lines, the one starting nearly from the base and leaving the costa beyond the middle, the other coming from the costa a little beyond it, and only slightly less oblique; at the angle where they meet are some silvery scales; above the acute angle which they form are three, white, wedge-shaped costal streaks, each wider than the other as they approach the apex, the third, pointing inward through the apical cilia, separated from the second by a blackish streak and ornamented at its outer end by two minute black dashes in the extremity of the apical cilia, which have thus a caudate appearance; the cilia below the bronzy brown pointed apex of the wing are also much ornamented, their base white in contrast with the narrow bronzy brown termen, some black spots lying along their middle beyond which their extremities are shining silvery grey; dorsal cilia dull greyish. *Exp. al.* 16 mm. *Hindwings* and cilia greyish brown, the latter tipped with whitish around the apex. *Abdomen* brownish cinereous. *Legs* whitish, the tarsi with four brownish spots.

Type ♂ (65395) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (65397) US. Nat. Mus.]

Hab. MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Three specimens.

The brown ground-colour of the forewings is distinctly more bronzy than in 5648 *acaciella* Busck, and the small white dash beneath the central white streak is absent.