

## 2. *Polyhymno convergens*, sp. n. (Tab. II. fig. 27.)

*Antennae* hoary white. *Palpi* hoary whitish, median joint slightly clothed beneath; terminal joint longer than median, shaded beneath. *Head* hoary white. *Thorax* hoary white, with two longitudinal greyish fuscous lines. *Forewings* greyish fuscous, longitudinally streaked with hoary whitish; a broad streak from the base below the costa extending over the fold, on the middle of which it is slightly broken; an arrow-like point following the fold towards the tornus; beyond the middle of the wing two hoary streaks, arising on the fold, converge towards the apex, nearly meeting the converging ends of four similar costal streaks of which the two outer are short and cuneiform, separated and followed by dark greyish fuscous, the tail-like apex also streaked above and beneath with fuscous and white alternately; an elongate silvery patch lies within the termen, the space around it and towards the costa and subapical incision fawn-brown; the cilia on the termen whitish, tipped with greyish fuscous, with three blackish spots along their base; the long cilia above the tornus hoary greyish. *Exp. al.* 16 mm. *Hindwings* brownish grey; cilia hoary grey. *Abdomen* hoary greyish, anal tuft ochreous. *Legs* hoary greyish; hind tibiae strongly hairy above, with two longitudinal streaks of greyish fuscous on their outer sides, tarsi banded with greyish fuscous, the first band much wider than the other four.

*Type* ♂ (65398) Mus. Wlsm. (*Godm-Salv. Coll.*) BM.

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII. (*H. H. Smith*). Unique.

Allied to 5648 *acaciella* Busck, but differing in the whitish lines converging towards the apex.

## 3. *Polyhymno colleta*, sp. n.

*Antennae* ivory-white, slightly clouded with fuscous beyond the basal third. *Palpi*, *Head*, and *Thorax* ivory-white, the latter with two slender brownish streaks along the middle. *Forewings* ivory-white, the dorsal third mottled with brown, which forms also a broken line along the fold, furcate near its outer end, the  $\sphericalangle$  running toward the apex; a rather broad brown band occurs along the costa, a slender white line running through it from before the middle of the costa to its outer and lower extremity; this is followed by a broader oblique white streak from the commencement of the costal cilia, nearly meeting the end of the slender white line below it; a pair of shorter, triangular, geminated streaks, the outer pair in the apical cilia separated by brown on the costa, the same colour running outward below them and forming a caudate apex in the cilia; preceding the termen and parallel to it are a few silvery metallic scales, the terminal cilia being ornamented by two black spots in a white patch, margined before by brown, and behind by brownish grey; dorsal cilia brownish grey. *Exp. al.* 11-12 mm. *Hindwings* pale bluish grey; cilia pale brownish grey. *Abdomen* leaden grey, whitish at the sides; anal tuft pale brownish ochreous. *Legs* ivory-white, with oblique brown streaks on the outer side of the tibiae, and brown bands across the tarsi.

*Type* ♂ (65807) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (65810) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Amula, 6000 ft., VIII-IX. (*H. H. Smith*). Four specimens.

## 4. *Polyhymno leucocras*, sp. n.

*Antennae* white at the base, shaded beyond. *Palpi* snow-white. *Head* and *Thorax* snow-white. *Forewings* snow-white, with a rather broad brown costal band, the dorsal space beneath the fold being also completely filled with brown, from which a brown streak is ejected outward in the direction of the apex; the white costal streaks are in the same positions as in the previous species (*colleta* Wlsm.), consisting of one long, slender, very oblique line commencing before the middle, followed by a shorter, less oblique line nearly meeting its apex, and two pairs of geminated white streaks in the costal and apical cilia; there is also a similar white patch in the terminal cilia, but instead of two black spots this contains a black