

the ends of the spoon-shaped claspers, which are rounded, but scarcely dilated posteriorly. *Legs* hairy throughout; pale cinereous, dusted with fuscous.

*Type* ♂ (66961) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (66963) US. Nat. Mus.]

*Hab.* MEXICO: GUERRERO: Tonalapa, VI. (*H. H. Smith*). Three specimens.

Veins 8 and 9 are stalked in the forewings.

### 50. *Acrolophus arcei* Druce.

*Anaphora arcéi* Druce Ann-Mag. NH. (7 s.) 7 442 (1901)<sup>1</sup>.

The antennae of the ♂ of this species are serrate, and minutely ciliate; the palpi overarch the thorax, and are roughly and distinctly tufted on each joint; the hind legs are hairy, except on the terminal joints of the tarsi. *Genitalia*: the uncus is somewhat angulate, double, with the points diverging, but the angle at the bend is rounded-off, not produced; the claspers are dilated posteriorly, inverted at the upper angle of their apex, excavate on their posterior margin, and produced into a somewhat broad lobe at the lower extremity.

*Type* ♂ (4550); ♀ (4551) Wlsm. Det. 1904) Mus. J. J. Joicey (*Witley*).

*Hab.* Central America<sup>1</sup>—MEXICO: VERA CRUZ: Jalapa, 4500 ft., 1887 (*W. Schaus*)—PANAMA<sup>1</sup>: CHIRIQUI (*Arcé*)<sup>1</sup>.

### 51. *Acrolophus uncispinis*, sp. n.

=\**Anaphora punctata* (p.) Druce Ann-Mag. NH. (7 s.) 7 443 (1901)<sup>1</sup>.

*Antennae* serrate, yellowish brown. *Palpi* overarching the thorax, thickly clothed throughout; brownish fuscous, the brush on the basal joint brownish ochreous. *Head* and *Thorax* brownish fuscous. *Forewings* with the ground-colour pale whitish ochreous, but almost completely suffused—except along and below the fold, and in a streak along the outer half of, and beyond, the cell—with fuscous, slightly tinged with purplish, on which are numerous small spots and strigulae of a still darker fuscous shade; the pale space below the fold is also somewhat sprinkled with these darker scales, and the plical patch is indicated by them on the fold, the other patches being obliterated and scarcely traceable; cilia pale dirty whitish ochreous, dusted with pale fuscous. *Exp. al.* 25–36 mm. *Hindwings* pale brownish fuscous, diluted to whitish ochreous about their base and on the cilia, through which runs a faint darker shade-line. *Abdomen* pale brownish fuscous: *genitalia*, uncus double, angular, the upper angle projecting to a length about equal with that of the lower point, and reaching as far as the ends of the claspers; these have their upper edge straight, but are dilated and rounded posteriorly beneath, terminating upward obtusely. *Legs* smooth; pale brownish cinereous.

*Type* ♂ (66984); ♀ (66985) *V. de Irazu*, Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PTT.* (6992–4, 67000, 67008–9) US. Nat. Mus.; (4498–9) Wlsm. Det. 1904) Mus. J. de Joannis; (\**punctata* Druce, *PT.*, 4556) Wlsm. Det. 1904) Mus. J. J. Joicey (*Witley*).]

*Hab.* Central America—GUATEMALA: SAN MARCOS: El Tumbador, 2500 ft., X–XI. 1880 (*G. C. Champion*)—NICARAGUA: Chontales (*E. M. Janson*)—COSTA RICA<sup>1</sup>: Volcan de Irazu, 6000–7000 ft. (*H. Rogers*); Candelaria Mts. (*C. A. Underwood*)<sup>1</sup>; San José (*Mus. Joannis*). Thirty-two specimens.

The ♀ is larger than the ♂, with very hairy palpi, which extend rather more than the length of the head beyond it. This species is extremely similar to *popeanella* Clms. and to *caprimulgus* Wlsm., but differs from both in the peculiar form of the uncus with its strongly projected upper angle.