

The Paratype of *Anaphora punctata* Druce, from Candelaria Mts. (*Underwood*)<sup>1</sup> (♂ 4556 Wlsm. Det. 1904), must be referred to *uncispinis* Wlsm.—Wlsm. MS. 812/1904.

## 52. *Acrolophus orasialis* Wkr.

*Palthis? orasialis* Wkr. Cat. Lp. BM. 19 886 (1859)<sup>1</sup>.

Type ♂ (1357) BM.

*Hab.* BRITISH HONDURAS [= \*HONDURAS<sup>1</sup> (*Miller*)<sup>1</sup>].

## 53. *Acrolophus caprimulgus*, sp. n.

*Antennae* serrate; brownish. *Palpi* overarching the thorax, densely clothed, tufted; brownish fuscous. *Head* and *Thorax* brownish fuscous. *Forewings* brownish ochreous, much suffused with dark brownish fuscous, except in a pale streak running along the fold, and in a shorter streak from the end of the cell, pointing toward the middle of the termen; the basal half of the costa and the whole of the cell very darkly suffused, and a dark excrescence from the central shade is thrown across the fold, forming the plical patch; on the outer half of the wing the pale ground-colour is slightly more apparent, showing three or four dark, confluent, costal streaks, the space below the fold being also shaded with fuscous; there is a slight admixture of dark ferruginous brown scales on the sites of the discal and plical patches, but the anteterminal patch is undistinguishable; cilia brownish cinereous, with a reduplicated dark shade-line. *Exp. al.* 26–32 mm. *Hindwings* pale fawn-brownish; cilia paler cinereous, with a shade-line running through them. *Abdomen* brownish cinereous, banded with dark brownish fuscous: *genitalia*, uncus double, short, angular, the angle not produced; claspers moderately wide, truncate, their slightly oblique posterior margin indented in the middle. *Legs* brownish cinereous, clothed with long, densely packed, hair-scales, except on the terminal joints of the tarsi.

Type ♂ (67043); ♀ (67044) *V. de Chiriqui*, Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PT. (67046) US. Nat. Mus.]

*Hab.* PANAMA: CHIRIQUI: Volcan de Chiriqui, 2000–3000 ft., 1881 (*G. C. Champion*): PANAMA: La Chorrera, IV–V. 1898 (*C. H. Dolby-Tyler*). Five specimens.

The ♀ has hairy erect palpi, rising scarcely above the base of the antennae.

## 54. *Acrolophus dentiger*, sp. n.

*Antennae* serrate, minutely ciliate; brownish fuscous. *Palpi* overarching the thorax, densely clothed and tufted; brownish cinereous, shaded with fuscous externally, the terminal joint sprinkled with pale-tipped scales. *Head* and *Thorax* brownish fuscous. *Forewings* pale brownish cinereous, evenly dusted and sprinkled with dark brownish fuscous, the usual discal, plical, and submarginal patches indicated by a thicker agglomeration of the same, the first discal being smeared nearly to the base; the fold, before and around the plical patch, shows the pale ground-colour scarcely obscured, and there is a somewhat clear streak of the same beyond the cell, but the amount of fuscous suffusion between the almost reticulate darker speckling is variable; the ultra-discal streak has a slight ochreous tinge; cilia brownish cinereous, with a median fuscous shade. *Exp. al.* 28–37 mm. *Hindwings* brownish fuscous; cilia pale brownish cinereous, with a median fuscous shade. *Abdomen* brownish fuscous: *genitalia*, uncus double, rather abruptly curved, but not angulate; claspers elongate, spoon-shaped, rounded posteriorly, with a strong inverted tooth about the middle of their lower edge. *Legs* brownish cinereous, with long closely laid scales.

Type ♂ (66978); ♀ (4631) *Jalapa*, Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [PT. (4629, 4632, 4634, 66980) US. Nat. Mus.]