

*Hab. Central America*¹—GUATEMALA: GUATEMALA: Guatemala City (*J. Rodriguez*)—COSTA RICA (*Van Patten*)¹; Volcan de Irazu, 6000–7000 ft. (*H. Rogers*)—PANAMA: CHIRIQUI: Volcan de Chiriqui, 2000–3000 ft., 1881–2 (*G. C. Champion*).

Anaphora punctata Druce was founded on two specimens: the Type (♂ 4557 Wlsm. Det. 1904) from Costa Rica (*Van Patten*), and a paratype (♂ 4556) from Candelaria Mts., Costa Rica (*Underwood*)—the latter must be referred to *uncispinis* Wlsm.

60. *Acrolophus pauper*, sp. n.

Antennae serrate, minutely ciliate; brownish. *Palpi* overarching the thorax, thickly clothed; brownish fuscous, the basal joint dark fuscous externally, and pale brownish cinereous in front on a level with the eyes. *Head* and *Thorax* brownish fuscous. *Forewings* brownish fuscous, somewhat paler, and with a rosy tinge, on and below the fold, except where interrupted by the large, triangular, dark brownish fuscous plical patch, and by a few minute dark upright dorsal strigulae; the discal patches are scarcely disconnected from the upper angles of the plical patch at either end, the one at the end of the cell obliquely placed, diffused and diluted nearly to the tornus; beyond it is an anteterminal series of about five dark fuscous spots, in an outwardly angulated line, a few indistinct spots lying along the termen at the base of the brownish fuscous cilia. *Exp. al.* 33 mm. *Hindwings* greyish brown; cilia brownish cinereous. *Abdomen* brownish cinereous: *genitalia*, uncus double, almost concave posteriorly through the projection, at its angle of deflection, where it is much widened; claspers of even width, with oblique hindmargin, deflected to a subobtuse apex, not reaching as far as the angle of the uncus. *Legs* densely, but not very roughly, clothed; brownish cinereous.

Type ♂ (66926) Mus. Wlsm. (*Godm-Salv. Coll.*) BM. [*PT.* (66928) US. Nat. Mus.]

Hab. MEXICO: VERA CRUZ: Cordova (*H. Kümeli*). Three specimens.

This species appears to be allied to *punctata* Druce, but differs in the somewhat less distinct dark blotches, in the detachment of the anteterminal spots from the outer discal patch, and, very noticeably, in the form of the uncus, projecting as it does beyond the claspers.

61. *Acrolophus gigantea* Druce.

Ankistrophorus giganteus Druce Ann-Mag. NH. (7 s.) 7 441 (1901)¹.

Antennae strongly serrate; pale brownish. *Palpi* recurved to behind the thorax; brownish fuscous, mottled with darker fuscous, especially on the outer side of the basal joint. *Head* and *Thorax* dark fuscous. *Forewings* dark greyish fuscous, with a lilac tinge, the costa narrowly touched with brownish throughout; there are a few dark transverse strigulae, and the usual dark patches are present, the plical indicated by the surrounding paler dorsal area; a small dark anteterminal patch lies in the same line as the larger one at the end of the cell; cilia brownish fuscous. *Exp. al.* 37–59 mm. *Hindwings* very dark umber-brown. *Abdomen* fuscous: *genitalia*, the uncus is angular, but, although the points are at right-angles with the stem, it is somewhat rounded-off at the bend, which projects as far as the ends of the stout widened claspers—these are straightened along their edges, both above and beneath, the upper edge slightly inverted posteriorly; their outer ends are almost straight, but project slightly on their middle. *Legs* greyish fuscous. (♂ 66916, *Atoyac*.) BM.

Type ♀ (*Orizaba*, 4527 Wlsm. Det. 1904) Mus. J. J. Joicey (*Witley*).

Hab. Central America—MEXICO¹: VERA CRUZ¹: Jalapa, 4500 ft., 1893 (*W. Schaus*); Atoyac (*H. H. Smith*; *Schumann*); Orizaba (*Boucard*)¹—COSTA RICA (Mus. J. de Joannis, 5000–1 Wlsm. Det. 1906)—PANAMA: CHIRIQUI: Bugaba, 800–1500 ft., 1881–3 (*G. C. Champion*).