

## ARISTOTELIA Hb. (pp. 22-6).

= *DORYPHORELLA* Ckrl.Type 5: *Anacamptis pulveratella* HS.*Xystophora* Hnmn-Wk. (1876) = *DORYPHORELLA* Ckrl. Ent. 21 163 (1888) nn. = § *DORYPHORA* Hnmn. (1870).7. *Aristotelia sarcodes* Wlsm. (p. 26)<sup>1</sup>.*Hab.* PANAMA<sup>1</sup>: CANAL ZONE<sup>1</sup>: Paraiso, II. 1911 (*A. Busck*).

## GALTICA Busck.

Type: *Galtica venosa* Busck (Busck 1914).*GALTICA* Busck Pr. US. Nat. Mus. 47 6-7 (1914).1. *Galtica venosa* Busck.*Galtica venosa* Busck Pr. US. Nat. Mus. 47 6-7 (1914)<sup>1</sup>.

Type ♂ (15817) US. Nat. Mus.

*Hab.* PANAMA<sup>1</sup>: Porto Bello, IV. (*A. Busck*)<sup>1</sup>.

## ANACAMPTIS Crt. (pp. 33-43).

24. *Anacamptis? veteranella* Z. (p. 43).

Three specimens received from Staudinger, in 1891, as *Tachyptilia veteranella* Z. (Mus. Wlsm. 6645-7) belong to the genus *Aristotelia*, but the shape of the wings is different from Zeller's figure of his species, and the markings do not agree.

25 (16.1). *Anacamptis peculella* Busck.*Anacamptis peculella* Busck Pr. US. Nat. Mus. 47 7 (1914)<sup>1</sup>.

Type ♂ (15818) US. Nat. Mus.

*Hab.* PANAMA<sup>1</sup>: Paraiso, Porto Bello, Trinidad River, and La Chorrera, II-V. (*A. Busck*)<sup>1</sup>.

26. *Anacamptis lagunculariella* Busck.

*Anacamptis lagunculariella* Busck Pr. US. Nat. Mus. 23 230-1 Pf. 1.6 (1900)<sup>1</sup>; Dyar Pr. Ent. Soc. Wash. 4 474 (1901)<sup>2</sup>; Busck Bull. US. Nat. Mus. 52 510 sp. 5706 (1902)<sup>3</sup>; Pr. US. Nat. Mus. 25 845, 848 (1903)<sup>4</sup>; 47 7 (1914)<sup>5</sup>.

Type ♂ (4937, *Palm Beach*) US. Nat. Mus.

*Hab.* AMERICA, N-C. United States<sup>1-5</sup>—FLORIDA<sup>1-5</sup>: Palm Beach, ⊕ *Laguncularia racemosa* (*H. G. Dyar*)<sup>1-2, 4-5</sup>. Central America<sup>5</sup>—PANAMA<sup>5</sup>: Corozal and La Chorrera, IV-V. (*A. Busck*)<sup>5</sup>.