

Sigmatomera flavipennis, O. Sack. Monogr. N. Am. Dipt. iii. (Addit. and Correct.), iv. p. 136.

Tipula nebulosa, Bellardi, Saggio &c. i. p. 6, t. 1. f. 4.—N.B. Is not a *Tipula*; seems to be allied to *Epiphragma*.

Limnophila undulata, Bellardi, Saggio &c. App. p. 3, t. 1. f. 2.—N.B. A true *Limnophila*.

Epiphragma solatrix, O. Sack. Monogr. &c. iv. p. 195.—Mexico, Orizaba (coll. Bellardi).

Limnobia stupens, Walk. Trans. Ent. Soc. Lond. n. ser. v. p. 332.—N.B. I suppose this is a *Gnophomyia*.

Tipula associans, Walk. Trans. Ent. Soc. Lond. n. ser. v. p. 332.

—— *dispellens*, Walk. l. c. p. 333.

—— *edwardsii*, Bellardi, Saggio &c. i. p. 8, t. 1. f. 2.

—— *quadrinaculata*, Bellardi, l. c. p. 9, t. 1. f. 3.

Pachyrrhina affinis, Bellardi, l. c. p. 10 (*Tipula*).

—— *mexicana*, Macq. Dipt. Ex. Suppl. i. p. 12.

TANYPREMNA, gen. nov.

Antennæ remarkably short, not much longer than the head, apparently 11-jointed. Legs very long; tarsi slender, whip-lash-shaped; venation of a *Tipula*, but anterior branch of the second vein still shorter than usual. Male forceps not incrassate, of a simple structure.

Tanypremna is a new genus which I propose for certain Tipulina related to *Dolichopeza*.

A detailed characterization will be published in a forthcoming paper of mine on Tipulidæ.

1. *Tanypremna opilio*, sp. n., ♂. (Tab. I. fig. 2, ♂*.)

Legs brown, the tibiæ whitish at the base, the tarsi white; thoracic dorsum brown; humeri and a portion of the pleuræ brownish-yellow; abdomen brown, the venter and male forceps brownish-yellow; wings slightly infuscated.

Length about 25 millim.

Hab. GUATEMALA, Purula in Vera Paz 5000 feet (*Champion*). A single male.

Head brownish-yellow; palpi and lips brownish; antennæ yellow, brownish towards the tips; a brown spot on the vertex. Collar brown, which colour is prolonged on the anterior part of the front coxæ; upper part of the mesonotum brown, the usual stripes being coalescent; a yellowish space in front of the suture, upon which two brown stripes are visible; a faint yellow line in the middle of the scutellum; metanotum brownish; humeri and pleuræ brownish-yellow, the latter with a brown stripe running from the dorso-pleural suture to the middle coxæ; sternum yellow. Coxæ and the base of the femora yellow, the rest of the femora and tibiæ dark brown, the latter have at the

* Through an error of the engraver, the seventh longitudinal vein and the last of the three veins issuing from the discal cell have been omitted from the figure.