

and not remarkably large; the wings, on the distal half, are almost imperceptibly tinged with greyish. About 6 millim. long.

The *Chrysopila apicalis*, v. d. Wulp, Tijdschr. &c. xxv. p. 119 (Guadeloupe), is still nearer; but I am by no means certain of the identity. Van der Wulp's name is pre-occupied by Rondani for a South-American species.

2. *Chrysopila* — ?

Hab. MEXICO, Northern Sonora (*Morrison*).

A pair (♂ ♀) from the above locality cannot be identified with any of the described species. Blackish-grey, with hyaline wings, yellowish tibiæ, black halteres, &c.

3. *Chrysopila* — ?

Hab. COSTA RICA, Irazu 6000 to 7000 feet (*Rogers*).

A pair (♂ ♀) from Irazu are certainly different from all the described species, but too badly preserved for description. The wings have a brown cross-band issuing from the darker brown stigma, crossing the middle of the first posterior cell, and reaching, somewhat attenuated, the cross-vein at the base of the fourth posterior cell; between this cross-band and the apex, which is likewise brown, there is a distinct hyaline cross-band. In our male specimen the brown is more spread out than in the female, and the hyaline band is cut off posteriorly by an infuscation of the last of the veins, issuing from the discal cell; in the female the brown markings are fainter and narrower, and therefore the hyaline cross-band broader, but less distinctly marked. Anal cell short, petiolate; a distinctly yellowish tinge at the proximal end of the stigma. Body brown in the female, with a golden-yellow pubescence, apparently forming double spots at the bases of the abdominal segments; I do not perceive any traces of this in the male. Halteres with a brown knob; legs brownish-yellow, the tarsi darker. Antennæ brown, the second joint reddish. Length 7–8 millim.

A third specimen (R. Sucio, Costa Rica, *Rogers*), a female, apparently belongs to the same species, but the brown markings on the wings are still fainter, and the brown cross-band almost obliterate.

4. *Chrysopila* — ?, ♂ ♀.

Hab. MEXICO, Ciudad in Durango 8100 feet (*Forrer*).

Two specimens. Body brown (♂), or greyish-brown (♀), with very scarce yellowish pile; antennæ dark brown; femora brown, the tibiæ brownish-yellow. Wings altogether hyaline, the stigma brown. Length 6–7 millim.