

feet, Venta de Zopilote 2000 feet, all in Guerrero (*H. H. Smith*), Temax in North Yucatan (*Gaumer*).—SOUTH AMERICA, Colombia<sup>2</sup>, Brazil<sup>1 3</sup>.

Ten specimens, all females. It is not at all improbable that the hitherto unknown male will be found to be identical with those described on p. 17 as *Syrphus*, no. 8.

The following species have been described by Bigot:—

*Ocyptamus infuscatus*, Bigot, Ann. Soc. Ent. Fr. 1883, p. 324.—Mexico.

—— *fraternus*, Bigot, loc. cit. p. 324.—Mexico.

—— *fenestratus*, Bigot, loc. cit. 1885, p. 251.—Mexico.

### BACCHA.

*Baccha*, Fabricius, Syst. Antl. p. 199 (1805).

#### *Synopsis of the recognized Central-American species.*

1. Alulæ small or rudimentary ; slender species (except *luctuosa*) 2.  
Alulæ of the usual size . . . . . 8.
2. Face in profile strongly projecting, forming nearly a straight  
edge with the front to the oral margin . . . . . *nasuta*, sp. n.  
Profile of face receding . . . . . 3.
3. Face in profile nearly straight and receding from the antennæ  
to the oral margin ; not at all tuberculate . . . . . 4.  
Face with a well-developed tubercle . . . . . 5.
4. Thorax metallic black ; small species . . . . . *gracilis*, sp. n.  
Thorax in great part red . . . . . *rubida*, sp. n.
5. Wings much shorter than the abdomen, hyaline, with a brown  
stigma ; front of female with fine transverse wrinkles ;  
abdomen very slender . . . . . *stenogaster*, Willist.  
Front of female without transverse wrinkles . . . . . 6.
6. Anal angle acuminate, with a concave basal border ; pleuræ  
and pectus yellow . . . . . *attenuata*, sp. n.  
Basal border of anal angle convex ; pectus, at least, metallic  
blue-black . . . . . 7.
7. Wings yellowish, with the anterior border indefinitely brown  
or brownish . . . . . *punctifrons*, sp. n.  
Wings uniformly deep yellowish-brown ; ocelli remote from  
vertex . . . . . *luctuosa*, Bigot?  
Wings hyaline, with the anterior border definitely dark  
brown . . . . . *laudabilis*, sp. n.