

8. Antennæ elongate, the first joint longer than the third; front tarsi of female flattened *conjuncta*, Wiedem.
- First antennal joint short 9.
9. Yellowish species; front with a more or less pronounced antenniferous projection, marked with a round black spot; front of female narrow above; mesonotum (except the sides) with its ground-colour black, but the black more or less concealed beneath ochraceous pollen; legs yellow, the hind femora or tibiæ more or less brownish or blackish 10.
- Species in great part black 11.
10. Black stripes of mesonotum conspicuous; very large species; wings brownish-yellow *phaeoptera*, Schiner.
- Black stripes of mesonotum inconspicuous, mostly concealed beneath light pollen; smaller species; wings more or less hyaline distally *livida*, Schiner.
11. Face, the four anterior tibiæ, and the hind metatarsi light yellow; wings brown along the costa; large species . . . *dolosa*, sp. n.
- Species not so marked 12.
12. Face projecting, yellow, with a black stripe; wings hyaline, with the subcostal (and costal, $\frac{2}{3}$) cells brown *clavata*, Fabr.
- Face not projecting 13.
13. Tip of the hind metatarsi and the following joints yellowish-white; wings strongly blackish throughout (σ), or with the distal part behind subhyaline (ρ) *lugubris*, sp. n.
- Distal joints of hind tarsi not yellowish-white 14.
14. Third and fourth abdominal segments each with two pairs of yellow dots, the anterior ones remote, the posterior pair approximate; legs wholly black; wings brown at the base, hyaline (σ) or subhyaline distally (ρ) *adspersa*, Fabr.
- Species not so marked 15.
15. Wings somewhat brownish in front; abdomen with yellow bands; mesonotum metallic bronze-green *ænea*, sp. n.
- Abdomen opaque and shining metallic, without yellow . . . 16.
16. Wings hyaline, with the subcostal cell brown; second and third abdominal segments with a single triangular opaque spot, the fourth and fifth segments with three spots . . . *concinna* (ρ), sp. n.
- Wings hyaline, with a narrow costal brown border reaching to the tip; second abdominal segment with a single spot, the third and fourth segments with two opaque spots . . . *cærulea*, sp. n.
- Wings brownish in front, subhyaline behind; abdominal segments opaque, with the margins shining; hind metatarsi blackish, the following joints yellow *concinna* (σ), sp. n.