

Hab. MEXICO (*Mus. Stockholm*)³.—ANTILLES^{1 2 4},

Stål alone records this species from Mexico; it is apparently common to the Antillean fauna.

3. *Tetyra farcta*. (Tab. III. fig. 1.)

Pachycoris farctus, Germ. Zeitschr. i. p. 92. 20¹; Walk. Cat. Het. Hem. i. p. 47.

Pachycoris guttipes, Walk. Cat. Het. Hem. i. p. 47².

Tetyra farcta, Stål, En. Hem. i. p. 5³.

Hab. MEXICO^{1 3} (*Mus. Berol.*), Orizaba, Oaxaca².—COLOMBIA³.

This species, as stated by Stål, is distinguished from the two preceding by the structure of the odoriferous orifices, "breviter ovalibus marginibus levissime elevatis, haud auriculatis." The specimen figured is the type form described by Walker as *P. guttipes*, in the collection of the British Museum.

PACHYCORIS.

Pachycoris, Burmeister, Handb. ii. 1, p. 391 (1835); Stål, Öfv. Vet.-Ak. Förh. 1867, p. 492.

This genus as now restricted is well represented in Central America; it is found in the West-Indian Islands, and extends southwards through Brazil and the Argentine Republic. Of the six species now associated as congeneric, two are here included. The remaining described forms are *P. fabricii*, Linn., *P. wilsonii*, Uhler (Cuba), *P. discrepans*, Uhler (N. Peru), and *P. obscuratus*, H.-S., described as from South America. Stål distinguishes this from the preceding genus in chief by "segmento sexto ventris feminarum apice medio late rotundato producto." The length of the head is greater than the breadth between the eyes.

1. *Pachycoris torridus*.

Cimex torridus, Scopoli, Ann. V. Hist.-Nat. p. 110. 109; Gmel. Syst. Nat. i. 4, p. 2134. 177.

Tetyra schousboei, Fab. Syst. Rhynch. p. 132. 18; Schjödte in Kröyer's Nat. Tidsskr. iv. p. 288. 8.

Pentatoma fabricii, Tigny, Hist. des Ins. ed. 2, iv. p. 290. f. 6.

Scutellera decorata, Perty, Del. An. p. 164, t. 33. f. 1.

Pachycoris fabricii, Burm. Handb. ii. 1, p. 392. 1; Hahn, Wanz. Ins. iii. p. 27. f. 249; Blanch. Hist. des Ins. p. 155. 3; Germ. Zeitschr. i. 1, p. 84. 5; A. et S. Hist. des Hém. p. 32. 2; Dall. List Hem. i. p. 31. 3, excl. syn. Linnæi et Fabricii; H.-S. Wanz. Ins. iv. p. 6. f. 349, excl. syn.

Pachycoris klugii, Germ. Zeitschr. i. 1, p. 85. 6; Dall. List Hem. i. p. 31. 2; H.-S. Wanz. Ins. iv. p. 5, f. 347 & 348; Burm. Handb. ii. 1, p. 392. 3¹.

Pachycoris linnæi, Hope, Cat. Hem. i. p. 12.

Pachycoris schousboei, Burm. Handb. ii. 1, p. 392. 2; Germ. Zeitschr. i. 1, p. 83. 4.

Pachycoris aquila, H.-S. Wanz. Ins. v. p. 7. f. 463.

Pachycoris stallii, Uhler, Proc. Ent. Soc. Phil. ii. p. 159.

Pæcilocoris æneiventris, Voll. Faune Ent. Ind. Neerl. i. p. 7, pl. 1. f. 3.

Pachycoris decoratus, Mayr, Reise der Freg. Novara, Hem. p. 25.

Pachycoris torridus, Stål, En. Hem. i. p. 5; Berg, Hem. Argent. p. 25.