

11. **Podisus congrex.** (Tab. I. fig. 20.)*Telepta congrex*, Stål, Stett. ent. Zeit. xxiii. p. 91. 33<sup>1</sup>.*Podisus congrex*, Stål, En. Hem. pt. 1, p. 51. 17.*Hab.* MEXICO<sup>1</sup> (*Mus. Berol.*); GUATEMALA, Capetillo, San Gerónimo (*Champion*); COSTA RICA, Irazu (*Rogers*).

The pronotal spines vary in colour from dark fuscous to bright rosy red. The size is also inconstant, specimens examined having been found to range from 9 millims. to 13 millims. in length. The specimen figured is from Costa Rica, and was compared and identified by the late Prof. Stål.

12. **Podisus invaria.** (Tab. IV. fig. 21.)*Arma invaria*, Walk. Cat. Het. i. p. 135. 15<sup>1</sup>.*Hab.* MEXICO, Oaxaca (*Brit. Mus.*).

The figure is from the type in the British Museum.

C. *Anterior lateral margins of the pronotum not crenulated.*

13. **Podisus iole.** (Tab. IV. fig. 1.)*Telepta iole*, Stål, Stett. ent. Zeit. xxiii. p. 92. 35<sup>1</sup>.*Podisus iole*, Stål, En. Hem. i. p. 52. 20.*Hab.* MEXICO (*coll. Sign.*).14. **Podisus thetis.** (Tab. IV. fig. 2.)*Telepta thetis*, Stål, Stett. ent. Zeit. xxiii. p. 92. 36<sup>1</sup>; En. Hem. i. p. 52. 21.*Hab.* MEXICO<sup>1</sup>; GUATEMALA, Zapote (*Champion*).

The specimen figured is typical, and from the collection of Dr. Signoret. A variety has been received from Guatemala which differs in the upper surface being dark olivaceous; the anterior half of the pronotal lateral borders luteous as in the typical form.

D. *Scutellum ornamented with a macular callosity at each basal angle.*

(*Subg.* *Tylospilus*, Stål.)

15. **Podisus acutissimus.** (Tab. II. fig. 22.)*Podisus (Tylospilus) acutissimus*, Stål, En. Hem. i. p. 53. 27<sup>1</sup>.*Tylospilus acutissimus*, Uhler, Bull. U.S. Geol. & Geog. Surv. ii. p. 283<sup>2</sup>.*Hab.* NORTH AMERICA, Colorado<sup>2</sup>, Texas<sup>1</sup>.—MEXICO<sup>1</sup>; GUATEMALA, San Gerónimo (*Champion*).

Stål enumerates the following varieties:—

Var. *a.*

Thorace posterius fusco vel nigricante.