

Pentatoma (*Mormidea*) *typhoeus*, Guér. in Sagra, Hist. de Cuba, Ins. p. 370³.

Oebalus typhoeus, Stål, Hem. Fabr. i. p. 27. 1; Townend Glover, Ills. Ins. Ord. Hem. t. 11. fig. 7.

Oebalus pugnax, Stål, En. Hem. ii. p. 22. 1⁴; Uhler, Bull. U.S. Geol. & Geogr. Surv. ii. p. 285⁵;

Proc. Bost. Soc. Nat. Hist. vol. xix. p. 377⁶.

Hab. NORTH AMERICA^{2 4}, Pennsylvania⁶, Maryland⁶, Virginia⁶, Carolina^{1 4 6}, Texas^{4 5}, Arizona⁵.—MEXICO^{4 5}; HONDURAS²; GUATEMALA, San Gerónimo (*Champion*).—CUBA^{3 5}; JAMAICA²; COLOMBIA^{4 5}.

Prof. Uhler states that this species "occurs at considerable altitudes on the Black Mountains of North Carolina, and is not unfrequent upon grassy and shrubby spots in Georgia and Florida." "It appears to lay eggs in late spring and towards the latter end of summer." "The Cuban form is usually larger, and has the lateral angles longer and more slender than in those from the United States."

SIBARIA.

Sibaria, Stål, En. Hem. ii. p. 23 (1872).

Stål thus differentiates the above:—"Hoc genus ad *Mormideam* appropinquat, sed differt capite antrorsum magis angustato, oculis majoribus, femoribus spinosis, tibiis distincte sulcatis, corii margine exteriori basi haud calloso; ab *Euchisto* et affinibus differt marginibus lateralibus thoracis obtusissimis, transversim rugosis." *Sibaria* at present includes but one species.

1. *Sibaria armata*. (Tab. V. fig. 17.)

Mormidea armata, Dall. List Hem. i. p. 215. 14¹; Stål, En. Hem. ii. p. 23. 1².

Hab. GUATEMALA, La Tinta, Vera Paz (*Champion*).—COLOMBIA²; BRAZIL^{1 2}.

A Guatemalan specimen is figured.

GALEDANTA.

Galedanta, Amyot et Serville, Hist. des Hém. p. 136 (1843); Stål, Öfv. Vet.-Ak. Förh. 1867, p. 527.

Only two species of *Galedanta* are at present recorded, both belonging to the Neotropical Region. It has considerable affinity with the following genus, *Euschistus*; but the lateral lobes of the head are longer than the central, and the scutellum possesses a tuberculous callosity in each basal angle.

1. *Galedanta myops*. (Tab. V. fig. 13.)

Cimex myops, Fabr. Syst. Rhynch. p. 155. 1.

Galedanta myops, Stål, Hem. Fabr. i. p. 26. 1; En. Hem. ii. p. 23. 1¹.

Hab. COSTA RICA, Cache (*Rogers*).—COLOMBIA (*coll. Dist.*); GUIANA¹.

BIOL. CENT.-AMER., Rhynch., October 1880.