

The large specimens with bluntly rounded angles have not yet been collected by me in company with the other form." These, agreeing with the *E. luridus*, Dall. (the form we have figured, with a black-coloured variety), have hitherto been received only from Guatemala; but the Berlin Museum contains both forms from Mexico. Prof. Uhler also states that in North America "it sometimes occurs in large numbers in late summer on bushes in damp situations." "A whitish bloom often covers the mature specimens in this region." This is also observable in the individuals of this species received from Guatemala.

3. *Euschistus rugifer*. (Tab. VI. figg. 8 & 9.)

Euschistus rugifer, Stål, En. Hem. ii. p. 26. 24¹.

Hab. MEXICO ¹.

The specimens figured are typical, and are in the collection of the Stockholm Museum. The species appears to be most variable in size.

4. *Euschistus bifibulus*.

Pentatoma bifibula, Palis. de Beauv. Ins. p. 148, Hém. pl. 10. fig. 5¹.

Euschistus pallipes, Dall. List Hem. i. p. 204. 8².

Euschistus bifibula, Dall. List Hem. i. p. 204. 9³; Stål, En. Hem. ii. p. 27. 25⁴; Stett. ent. Zeit. xxiii. p. 100. 58⁵.

Hab. MEXICO ^{4 5} (*coll. Signoret*), Oaxaca (*coll. Dist.*); HONDURAS ^{3 4}; GUATEMALA, San Gerónimo (*Champion*); COSTA RICA (*Van Patten*).—CUBA ⁴; SAN DOMINGO ¹; JAMAICA ²; COLOMBIA ⁴.

5. *Euschistus crenator*.

Cimex crenator, Fabr. Ent. Syst. iv. p. 101. 87; Syst. Rhynch. p. 162. 38.

Pentatoma obscura, Palis. Beauv. Ins. p. 149, Hém. pl. 10. figg. 7 & 9.

Pentatoma pustulata, Palis. Beauv. Ins. p. 185, Hém. pl. 11. fig. 2.

Euschistus obscurus, Dall. List Hem. i. p. 208. 19¹; Stål, Stett. ent. Zeit. xxiii. p. 100. 59².

Pentatoma (Mormidea) obscura, Guér. in Sagra, Hist. de Cuba, Ins. p. 366³.

Pentatoma (Mormidea) pustulata, Guér. in Sagra, Hist. de Cuba, Ins. p. 368⁴.

Euschistus crenator, Stål, Hem. Fabr. i. p. 26. 2; En. Hem. ii. p. 27. 26⁵; Uhler, Bull. U.S. Geol. & Geogr. Surv. ii. p. 286. 5⁶.

Hab. NORTH AMERICA ⁶, Texas ^{5 6}, Florida ^{5 6}.—MEXICO ^{2 5 6} (*Mus. Berol.*); CENTRAL AMERICA ¹; COSTA RICA (*Van Patten*), Irazu (*Rogers*), Panama (*coll. Dist.*), Matachin (*Thiene, coll. Oberthür*).—CUBA ^{3 4 6}; COLOMBIA ⁵.

6. *Euchistus spurculus*. (Tab. VI. fig. 10.)

Euschistus (Lycipta) spurculus, Stål, Stett. ent. Zeit. xxiii. p. 100. 58¹; En. Hem. ii. p. 28. 30².

Hab. MEXICO ^{1 2}; GUATEMALA, Zapote (*Champion*).