

- divergent, subequally reticulated, obliquely unifasciate, the subcostal area biseriate behind; the margins of the elytra and the pronotum pilose; head without spines *vesiculosa*, n. sp.
- b.* Antennæ with joint 1 about four times as long as 2 *. Pronotum with the membranous margins moderately wide, the median carina foliaceous, and the hood rather small; elytra long and unequally reticulated, the subcostal area biseriate; pronotum pilose in front; head with five spines *longipennis*, n. sp.
- c.* Antennæ with joint 1 three or three and a half times as long as 2. Head with from three to five spines.
- a'.* Median carina of the pronotum strongly foliaceous; subcostal area of the elytra bi- or uniseriate.
- a''.* The membranous margins of the pronotum broadly, arcuately dilated, with three or four rows of areolæ; hood very large.
- a'''.* Outer carinæ of the pronotum obsolete; elytra widely and unequally reticulated, very obliquely unifasciate; pronotal and elytral margins without distinct setæ *tumida*, n. sp.
- b'''.* Outer carinæ of the pronotum moderately foliaceous; elytra unequally reticulated, obliquely unifasciate; pronotal and elytral margins setose *setigera*, n. sp.
- b''.* The membranous margins of the pronotum moderately wide, with two rows of areolæ; hood rather small.
- c'''.* Elytra widely and unequally reticulated, not fasciate; pronotum and elytral margins set with long fine hairs *fimbriata*, n. sp.
- d'''.* Elytra very widely and unequally reticulated, obliquely unifasciate; pronotum and elytral margins indistinctly pilose *fuscofasciata*, n. sp.
- b'.* Median carina of the pronotum feebly foliaceous, sometimes more raised than the outer ones.
- c''.* The membranous margins of the pronotum broadly and subangularly dilated, closely reticulated, the hood large; elytra obliquely unifasciate, the costal area with about four rows of areolæ, the subcostal area triseriate *elata*, n. sp.
- d''.* The membranous margins of the pronotum moderately wide, with two rows of areolæ, the hood small; costal area of the elytra narrower, with two rows of areolæ, the subcostal area biseriate.
- e'''.* Elytra narrowly and very obliquely unifasciate, the reticulation wide and unequal, the apical margin oblique *lineata*, n. sp.
- f'''.* Elytra obliquely and faintly unifasciate, the reticulation unequal, the apical margin rounded *gracilentata*, n. sp.
- g'''.* Elytra longer and not fasciate, but with the nervures darker on the apical third or more, the apical margin rounded.
- a⁴.* Antennæ and legs very elongate; elytra widening behind *angustata*, n. sp.
- b⁴.* Antennæ and legs moderately elongate; elytra (when closed) narrowing behind *tenuis*, n. sp.

* Foreshortened in our figure.