

in the costal area infusate, the areolæ of the latter hyaline, the tarsi usually fuscous at the tip. Head with slender, porrect, frontal spines; antennæ very shortly pilose, reaching to the basal third of the elytra, stout, extremely shortly pilose, joint 2 slightly shorter than 1, 3 nearly four times as long as 4, 4 longer than 1 and 2 united. Pronotum tricarinate, the median carina raised and projecting in front, the marginal carinæ prominent, the interspaces coarsely, closely punctate, reticulate behind. Elytra long, somewhat oval in shape, rounded at the tip, the costal margin slightly sinuate at about one-third from the apex; discoidal area widely reticulated, the nervures surrounding it prominent, the outer one curved behind; subcostal area rather broad, biseriate; costal area prominent to the base, uniseriate to beyond the middle, biseriate towards the apex, the areolæ rather large. Rostral groove subparallel, the rostrum extending to a little beyond the metasternum.

Length  $4-4\frac{1}{2}$ , breadth  $1\frac{1}{2}-1\frac{3}{5}$  millim. ( $\sigma$  ♀.)

*Hab.* BRITISH HONDURAS, R. Hondo (*Blancaneaux*); GUATEMALA, Chiacam, San Juan, and San Gerónimo in Vera Paz (*Champion*).

Eight examples. Differs from *Teleonemia* proper in having two rows of areolæ in the costal area of the elytra towards the apex. The antennæ are formed very much as in *T. validicornis* and *T. forticornis*. A specimen from San Gerónimo is figured.

## 12. *Teleonemia variegata*, n. sp. (Tab. III. figg. 15; 15 a, profile.)

Elongate, opaque; brownish-testaceous or testaceous, fuscous beneath, the elytra mottled with blackish or fuscous, and with some of the transverse nervures in the costal area black, the areolæ of the latter hyaline, the apical joint of the antennæ partly or entirely fuscous, the tarsi black, the femora and tibiæ sometimes slightly annulated with fuscous. Head with slender porrect frontal spines; antennæ long, moderately stout, very slightly pilose, joint 2 slightly shorter than 1, 3 three or three and one-half times longer than 4, 4 a little longer than 1 and 2 united. Pronotum truncate in front, tricarinate, the median carina not raised anteriorly, the marginal carinæ moderately prominent, the interspaces coarsely, closely punctate, reticulate behind. Elytra long, somewhat oval in shape, constricted at about one-third from the apex, the apices rounded; discoidal area widely reticulated; the areolæ not very deeply impressed, the nervures surrounding it prominent, the outer one curved behind; subcostal area biseriate; costal area prominent to the base, uniseriate to beyond the middle, irregularly biseriate for a short distance towards the apex, the areolæ rather large. Rostral groove gradually widening, the rostrum extending to the second ventral suture.

Length  $4\frac{1}{5}-4\frac{1}{2}$ , breadth  $1\frac{1}{2}-1\frac{7}{8}$  millim.

*Hab.* MEXICO, Omilteme in Guerrero 8000 feet (*H. H. Smith*); GUATEMALA, Capetillo 4000 feet (*Champion*).

Two specimens from each locality. Near *T. cylindricornis*, but with the antennæ shorter and not so stout, the median carina of the pronotum neither raised nor projecting in front, the rostrum longer, the costal area of the elytra with only a few additional areolæ near the sinus, &c. A Mexican example is figured.

## 13. *Teleonemia picta*, n. sp. (Tab. III. figg. 16; 16 a, profile.)

Elongate, opaque; testaceous or ochraceous, the disc of the pronotum, the carinæ excepted, the inner part of the discoidal area, a transverse ante-median fascia between it and the costal margin, widening inwards, the nervures at the costal sinus, and a large apical patch, extending obliquely forwards to the base of the sutural area, more or less fuscous; the body beneath piceous; the antennæ fuscous or fusco-ferruginous, the apical joint black; the legs ferruginous or obscure ferruginous, with the tarsi blackish. Head with a rather long decumbent frontal spine, the two short ones below it approximating at the tip; antennæ moderately stout, about as long as the elytra, joint 2 slightly shorter than 1, 3 three times as long as 4, 4 longer than 1 and 2 united. Pronotum sharply tricarinate, the outer carinæ