

*d'''*. Joint 3 not or not much longer than 2.

*a*<sup>4</sup>. Form narrow, elongate; lateral angles of the pronotum emarginate; connexivum (♂) not visible from above . . . . *attenuatus*, n. sp.

*b*<sup>4</sup>. Form comparatively short; lateral angles of the pronotum obliquely truncate or feebly emarginate; connexivum (♂ ♀) broad.

*a*<sup>5</sup>. Lateral angles of the pronotum dilated and reflexed on their anterior edge; pronotum and scutellum finely punctured. *inæqualis*, n. sp.

*b*<sup>5</sup>. Lateral angles of the pronotum simple in front; pronotum and base of scutellum coarsely punctured.

*a*<sup>6</sup>. Anterior lobe of the pronotum coarsely granulate, more or less spiculate at the sides, infusate in the male, the posterior portion feebly convex or somewhat flattened; scutellum moderately depressed below the base, in the male black, usually with two yellow spots on each side externally . . . . . *stali*, Handl.

*b*<sup>6</sup>. Anterior lobe of the pronotum more finely granulate.

*a*<sup>7</sup>. Pronotum convex behind, the anterior lobe flavous in the male; scutellum strongly depressed below the base, in the male with the base, a median fascia, and an apical spot more or less fuscous . . . . . *lepidus*, Stål.

*b*<sup>7</sup>. Pronotum flattened behind; scutellum feebly depressed below the base. [The male with the anterior and posterior lobes of the pronotum, the sides excepted, and a broad median vitta on the scutellum, fuscous.] . *prehensilis*, F.

*d'*. Head and anterior half of the pronotum with setiferous spiculæ, the rest of the surface conspicuously granulate; form short and broad . . . . . *spiculosus*, n. sp.

### 1. *Macrocephalus notatus*. (Tab. IV. fig. 10, ♂.)

*Macrocephalus notatus*, Westw. Trans. Ent. Soc. Lond. iii. p. 24 (1841)<sup>1</sup>; Handl. Ann. k. k. naturhist.

Hofmus. xii. p. 188, fig. 22 (scutellum), t. 9. figg. 27, 28 (antenna)<sup>2</sup>.

*Macrocephalus incisus*, Stål, Stett. ent. Zeit. 1862, p. 440 (♂)<sup>3</sup>; Enum. Hemipt. v. p. 135<sup>4</sup>.

*Macrocephalus cliens*, Stål, Stett. ent. Zeit. 1862, p. 440 (♂)<sup>5</sup>.

*Hab.* MEXICO<sup>2-5</sup>, Pinos Altos in Chihuahua (*Buchan-Hepburn*), Teapa in Tabasco (*H. H. Smith*), Tabasco (*Mus. Holm.*<sup>2</sup>), Temax in N. Yucatan (*Gaumer*); BRITISH HONDURAS, Belize (*Blancaneaux*); GUATEMALA<sup>2</sup>, Chiacam, San Juan, Panzos, Chacoj, and La Tinta in Vera Paz, El Reposo, Pantaleon, Mirandilla (*Champion*), Escuintla (*Mus. Vind. Cæs.*); COSTA RICA<sup>2</sup>; PANAMA, Veraguas (*Mus. Berol.*<sup>2</sup>). — COLOMBIA<sup>1 2</sup>; VENEZUELA<sup>2</sup>.

*Var.* The antero-lateral tooth of the pronotum less prominent; the scutellar callus smaller and shorter, not reaching the middle (Handl. *loc. cit.* fig. 24). (♂ ♀.)

*Hab.* GUATEMALA, San Gerónimo (*Champion*); COSTA RICA<sup>2</sup>.