

1. *Lyctocoris campestris*.

Acanthia campestris, Fabr. Ent. Syst. iv. p. 75¹.

Lyctocoris campestris, Reut. Öfv. Vet.-Ak. Förh. 1871, p. 409²; Monogr. Anthocorid. p. 7³.

Lyctocoris fitchii, Reut. Öfv. Vet.-Ak. Förh. 1871, p. 557⁴; Stål, Enum. Hemipt. iii. p. 101⁵.

Hab. NORTH AMERICA, New York^{4 5} to Texas³.—MEXICO, Chiapas (*Richardson*).—EUROPE³; ASIA MINOR³; NEW ZEALAND³ &c.

A specimen of this insect has been found by us amongst a collection of bird-skins received from Chiapas. For the rest of the synonymy, see Dr. Reuter's Monograph.

LASIOCHILUS.

Lasiochilus, Reuter, Öfv. Vet.-Ak. Förh. 1871, p. 562; Monogr. Anthocorid. pp. 5, 13; Stål, Enum. Hemipt. iii. p. 102.

Dilasia, Reuter, Öfv. Vet.-Ak. Förh. 1871, p. 563.

Hapa, Buchanan White, P. Z. S. 1878, p. 465.

Subgen. *Semiotoscelis*, Reuter, Monogr. Anthocorid. pp. 15, 24.

Of the sixteen described species of this widely-distributed genus*, ten are American. All but one of the Central-American forms appear to be new.

Its chief characters are the short, backwardly-curved orifice of the metastethium and the strongly pilose elytra, the outer margins of the embolium appearing ciliate.

a. Pronotum with a median fovea or short longitudinal sulcus on the anterior lobe behind.

a'. Antennæ with joints 1 and 2 moderately thickened.

a''. Pronotum much narrowed anteriorly, more than twice as wide at the base as at the apex; elytra closely punctured *punctipennis*, n. sp.

b''. Pronotum moderately narrowed anteriorly, not twice as wide at the base as at the apex.

a'''. Elytra closely and finely punctured, fuscous *microps*, n. sp.

b'''. Elytra sparsely and finely punctured, maculate *reuteri*, n. sp.

c'''. Elytra with the clavus coarsely and closely, and the other parts finely and very sparsely, punctured, testaceous *pallidulus*, Reut.

b'. Antennæ with joints 1 and 2 stout; pronotal fovea deep; elytra with the clavus coarsely, and the other parts more finely, punctured, fuscous, with the shoulders pale *foveicollis*, n. sp.

b. Pronotum with the anterior lobe sulcate down the middle from the ante-apical groove to the base; elytra with the clavus coarsely, and the other parts more finely, punctured, testaceous.

c'. Pronotum moderately narrowed anteriorly, nearly twice as wide at the base as at the apex: body ovate *sulcatus*, n. sp.

d'. Pronotum broad in front, about one-third wider at the base than at the apex, the sulcus very deep: body elongate *divisus*, n. sp.

* *L. nebulosus* and *L. pictus*, Uhler, belong to *Asthenidea*.