

This species exhibits little variation; and the two specimens described by Walker under the above names are identical. The habitats of these were unrecorded.

4. **Fidicina spinicosta.** (Tab. II. figg. 15, 15 *a*, 15 *b*.)

*Cicada spinicosta*, Walk. List Hom. i. p. 126. 64<sup>1</sup>.

*Cicada lacrines*, Walk. ibid. p. 132. 70; Stål, Öfv. Vet.-Ak. Förh. 1862, p. 481.

*Fidicina lacrines*, Stål, ibid. p. 485.

*Hab.* COSTA RICA (*Mus. Berol.*); PANAMA (*coll. Dist.*).—GUIANA, Cayenne (*coll. Sign.*); BRAZIL, Para<sup>1</sup>.

Walker, under the name of *C. spinicosta*, subsequently incorporated specimens of his other species, *C. semilata*; and Stål necessarily thus wrote *C. lacrines*=*C. spinicosta* ("ad partem"). There can, however, be no doubt as to the type, which I have examined in the British Museum, and which agrees with the Costa-Rican specimen here figured contained in the collection of the Berlin Museum.

5. **Fidicina semilata.** (Tab. III. figg. 7, 7 *a*, 7 *b*.)

*Cicada semilata*, Walk. List Hom. i. p. 122. 59<sup>1</sup>.

*Cicada passer*, Walk. ibid. p. 124. 61<sup>2</sup>; Stål, Öfv. Vet.-Ak. Förh. 1862, p. 481.

*Cicada brizo*, Walk. ibid. p. 125. 63.

*Cicada melisa*, Walk. ibid. p. 127. 65<sup>3</sup>.

*Cicada melina*, Walk. ibid. p. 128. 66.

*Cicada panyases*, Walk. ibid. p. 133. 71<sup>4</sup>.

*Cicada pidytes*, Walk. ibid. p. 134. 72.

*Cicada physcoa*, Walk. ibid. p. 135. 73.

*Cicada braure*, Walk. ibid. p. 136. 74<sup>5</sup>.

*Cicada solemnis*, Walk. ibid. p. 143. 81<sup>6</sup>.

*Fidicina semilata*, Stål, Öfv. Vet.-Ak. Förh. 1862, p. 485.

*Hab.* COSTA RICA (*van Patten*; *coll. Dist.*).—ST. LUCIA<sup>16</sup>; VENEZUELA<sup>23</sup>; GUIANA, Demerara<sup>4</sup>, Cayenne<sup>1</sup>; TRINIDAD (*coll. Dist.*); BRAZIL, Para<sup>5</sup>, Ega (*de Mathan*; *coll. Oberthür*).

This species is not of an exceedingly variable nature, as the above synonymy would apparently indicate, but is rather an example of the late Mr. Walker's treatment of the subject. It is allied to *F. spinicosta*, but differs by the longer and less robust body, the more elongate and less tumid face, shape of the opercula, &c.

6. **Fidicina fumea**, n. sp. (Tab. IV. figg. 4, 4 *a*, 4 *b*.)\*

♀. Head and thorax above olivaceous. Head with the front marked with a central lunate spot, a central longitudinal line and the basal margin black; vertex broadly black between the eyes, which are bronzy;

---

\* On the preceding page it was stated that five species of *Fidicina* were enumerated. The present (sixth) species was received from Mr. Champion since that statement was printed.