

1. **Coanaco guttata.** (Tab. IV. figg. 10, 10 *a*, 10 *b*.)*Enchophora guttata*, White, Ann. & Mag. Nat. Hist. 1846, xvii. p. 331 ¹.*Aphæna guttata*, Walk. List Hom. ii. p. 276. 14.*Hab.* MEXICO, Jalapa (*Höge*); GUATEMALA, San Isidro (*Champion*); PANAMA, Bugaba, Tolé (*Champion*).

White gave the somewhat vague locality "South America" ¹, a habitat in any case too loose for any scientific purpose, and, in this instance, possibly erroneous.

2. **Coanaco ornanda**, n. sp. (Tab. IV. figg. 13, 13 *a*, 13 *b*.)

Head, pronotum, and mesonotum fuscous. Tegmina fuscous, the claval area much paler; the apical area ornamented with pale spots, and the basal area with scattered black spots, some of which have red centres. Wings dark fuscous, with about two thirds of the area from the base ornamented with bluish-grey spots, and with a few minute greyish spots at the apex. Abdomen above dark sanguineous, the two basal segments, some median segmental spots, and the anal appendage black. Body beneath and the legs dull reddish, the face and the margins of the prosternum dull luteous.

Long. excl. tegm. 16 millim., exp. tegm. 45 to 48 millim.

Hab. PANAMA, Chiriqui (*Ribbe*, *Mus. Vind. Cæs.*), Bugaba, Tolé (*Champion*).

OBIA, gen. nov.

The face in this genus is distinctly longer than broad, the outer margin carinate, slightly sinuate at about the middle of the lateral margins, and moderately amplified posteriorly; it has three central subparallel carinæ. The arrangement of the claval veins is as in *Aphana* and *Amantia*.

1. **Obia tenebrosa**, n. sp. (Tab. V. figg. 17, 17 *a*.)

Head, pronotum, and mesonotum ochraceous-brown; the pronotum with a black cruciform impression, and the mesonotum with blackish spots and markings. Tegmina reddish-ochraceous, the basal area with blackish markings, the apical area duller in hue and marked with large irregular greyish spots. Wings sanguineous, with the apical, posterior, and abdominal margins broadly blackish. Abdomen above blackish, with the lateral margins brownish-ochraceous; head beneath and the sternum and legs, pale brownish-ochraceous, more or less spotted and marked with blackish; abdomen beneath blackish, the tibiæ somewhat distinctly annulated with that colour.

Long. excl. tegm. 15 millim., exp. tegm. 46 millim.

Hab. MEXICO, Jalapa (*coll. Dist.*).

AMALIVACA, gen. nov.

In this genus the face is longer than broad, and its lateral margins are amplified and sublunate posteriorly; it has two central carinæ united before the apex, and the surface between the carinæ is concavely excavated; the facial base is angularly deflexed.