

## 2. *Poblicia constellata*. (Tab. IV. figg. 6, 6 a.)

*Poiocera constellata*, Walk. List Hom., Suppl. p. 50<sup>1</sup>.

*Poiocera tricolor*, Gerst. in Wieg. Archiv f. Naturg. xxvi. p. 238, t. 11. f. 11<sup>2</sup>.

*Hab.* MEXICO<sup>1 2</sup>, Jalapa (*Höge*); GUATEMALA, San Gerónimo, San Joaquin in Vera Paz (*Champion*).

The typical specimen described by Gerstaecker under the name of *P. tricolor* is here figured.

## 3. *Poblicia atomaria*. (Tab. V. figg. 2, 2 a.)

*Poiocera atomaria*, Walk. List Hom., Suppl. p. 49<sup>1</sup>.

*Hab.* MEXICO<sup>1</sup>; GUATEMALA, Calderas (*Champion*).

This name is not a synonym of *P. constellata*, as stated by Stål (Æfv. Vet.-Ak. Förh. 1862, p. 486), who was evidently led into the error by only examining the "unset" specimens of the species in the British-Museum collection.

## DOMITIA.

*Domitia*, Stål, Hem. Afr. iv. p. 138 (1866); Berl. ent. Zeitschr. 1866, p. 390; Stett. ent. Zeit. xxxi. p. 290 (1870).

I place the following species in this genus, of which I fortunately possess specimens of the type as given by Stål, viz. *D. basistella*, Walk., an Antillean insect. With this type our species generally agree, excepting in the form of the carinal markings of the face, a character alone too insignificant for fresh generic subdivision.

## 1. *Domitia neotropicalis*, n. sp. (Tab. V. figg. 3, 3 a.)

Head and eyes ochraceous; pronotum olivaceous-green, with discal black spots; mesonotum olivaceous-green, tinged with ochraceous, with four central longitudinal and a marginal series of rounded black spots; abdomen dull sanguineous, the base and a series of transverse discal streaks black; the posterior segmental margins narrowly pale greenish-ochraceous. Tegmina fuscous, with the veins and reticulations green spotted with ochraceous; the costal spots very distinct; apical area pale fuscous-hyaline; the venation fuscous. Wings pale fuscous-hyaline, the venation fuscous; the basal third black, with pale bluish-green reticulations and some basal spots of the same colour. Face and sternum pale olivaceous, shaded with brown; legs brownish-olivaceous. Abdomen beneath as above, but duller in hue. Rostrum reaching the penultimate segment.

Long. excl. tegm. 16 to 19 millim., exp. tegm. 54 to 58 millim.

*Hab.* MEXICO, Jalapa, environs of city (*Höge*); GUATEMALA, San Gerónimo (*Champion*); NICARAGUA, Chontales (*Janson*); PANAMA, David (*Champion*).

This species is allied to *D. flavo-punctata* (Perty), from which it differs by the greater hyaline area of the wings, different colour of the abdomen, &c.