

This is a very well-marked species, the length and the black linear markings of the prolonged head rendering it easily recognizable.

**2. Dictyophara ferocula, n. sp.** (Tab. VI. fig. 2.)

Head very long, about equal in length to that of the intermediate tibiae. The prolonged head is pale castaneous, the carinae greenish; pronotum and mesonotum pale obscure castaneous, with the carinae greenish. Tegmina pale hyaline, pale greenish at the base, the venation pale fuscous. Legs pale brownish, the extreme apices of the tibiae and tarsi a little darker.

Long. excl. tegm. 18 millim., long. head 7 millim., exp. tegm. 32 millim.

*Hab.* PANAMA, Matachin (*coll. Dist.*).

The great length of the head is sufficient to adequately separate this species from any of the others here enumerated.

**3. Dictyophara nodivena.** (Tab. VI. fig. 3.)

*Dictyophara nodivena*, Walk. Insect. Saundersiana, p. 37<sup>1</sup>.

*Hab.* PANAMA, Bugaba, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).—BRAZIL, Pará<sup>1</sup>.

**4. Dictyophara curviceps.** (Tab. VI. fig. 4.)

*Nersia curviceps*, Stål, Bidr. Rio Jan. Hem. p. 64<sup>1</sup>; Stett. ent. Zeit. xxv. p. 50. 353<sup>2</sup>.

*Dictyophara dioxys*, Walk. List Hom., Suppl. p. 61<sup>3</sup>.

*Hab.* MEXICO<sup>1 2 3</sup>; GUATEMALA, Zapote, Panzos in Vera Paz (*Champion*).

Stål's publication is dated January 1858, and Walker's List bears the date March 1858, so Stål's name seems clearly to have priority.

**5. Dictyophara brachyrhina.** (Tab. VI. fig. 5.)

*Dictyophara brachyrhina*, Walk. List Hom. ii. p. 317. 31<sup>1</sup>.

*Hab.* GUATEMALA, El Reposo, Zapote (*Champion*).—COLOMBIA<sup>1</sup>.

**6. Dictyophara obtusifrons.** (Tab. VI. fig. 6.)

*Dictyophara obtusifrons*, Walk. List Hom. ii. p. 318<sup>1</sup>.

*Hab.* PANAMA, Bugaba, Volcan de Chiriqui 2500 to 4000 feet, David (*Champion*).—GUIANA; AMAZONS; ANTILLES (*coll. Dist.*).

This insect is allied to *D. truncata*, Walk., a species recorded from Pará, and of which I possess a good series from Demerara and the Antilles. It principally differs from *D. truncata* in the shorter and more quadrate prolongation of the head. No locality is affixed to Walker's description<sup>1</sup>.

**7. Dictyophara herbida.** (Tab. VI. fig. 7.)

*Dichoptera herbida*, Walk. List Hom. ii. p. 306<sup>1</sup>.

*Dictyophara inexacta*, Walk. Ins. Saundersiana, p. 38<sup>2</sup>.