

2. **Flata rotundior**, sp. n. (Tab. VII. figg. 11, 11 a.)

Præcedenti affinis, sed dilutius virescens, et margine costali tegminum magis rotundato, angulo exteriori toto sensim rotundato, et margine apicali obliquiori facile distinguenda.

Closely allied to the preceding, but of a lighter green colour, with the tegmina (viewed from the side) distinctly narrower, and the costal margin more gradually rounded, the exterior angle being entirely and gradually rounded off, and the apical margin somewhat oblique; the reticulation, moreover, is coarser; the head not quite so much produced, but the difference in this character is very slight.

Long. 16-17 millim.; lat. ad hum. 3-4 millim.

Hab. MEXICO, Orizaba (*Bilimek, in Mus. Vind. Cæs.*); GUATEMALA, Panima and Senahu in Vera Paz (*Champion*).

I have found one specimen of this distinct species among the unnamed undetermined material belonging to the Vienna Museum. We figure a specimen from Panima.

3. **Flata punctata**, sp. n.

Præcedenti affinis, sed minor, marginibus tegminum magis parallelis, margine interiori recto, angulo apicali obtuso haud reflexo, margine exteriori fere recto, angulo apicali obtuse rotundato, margine apicali levissime rotundato; tegminibus subtiliter reticulatis.

Allied to *F. rotundior*, but smaller, with the margins of the tegmina somewhat parallel, interior margin straight, with the inner apical angle obtuse (not reflexed and acute, as in the two preceding species), exterior margin nearly straight, with the outer apical angle obtusely rounded, apical margin very slightly rounded; reticulation very fine.

Long. 13 millim.; lat. ad hum. 3 millim.

Hab. MEXICO (*Bilimek, in Mus. Vind. Cæs.*).

The single specimen above described stands in the Vienna Museum under the name of *Phyllyphanta punctata*, Signoret; this, however, as is the case with so many species of Signoret, is simply a manuscript name, for the species does not appear to have been described. It is very distinct.

4. **Flata rufocinctata**, sp. n. (Tab. VII. figg. 12, 12 a.)

Prasina, nitidiuscula, marginibus tegminum peranguste rufo-limbatis; capite brevi, perobtuse triangulariter producto; pronoto ad medium carinato; scutello tricarinato; tegminibus haud latis, margine exteriori leviter obliquo, angulo apicali obtuse rotundato, margine interiori recto, angulo apicali recto, margine apicali truncato; pedibus testaceis; tarsis tibiisque rufis.

Green, with the margins of the tegmina very narrowly bordered with red, the colour almost ceasing towards the base on the costal margin; head short, obtusely, triangularly produced; pronotum about as long as the head, with a central carina; scutellum tricarinate; tegmina comparatively narrow, exterior margin slightly oblique, with the apical angle obtusely rounded, interior margin straight, with the apical angle a sharp right angle, almost acute, apical margin straight, truncate; reticulation fine and very distinct; legs mostly red.

Long. 12 millim.; lat. ad hum. $2\frac{1}{2}$ millim.

Hab. PANAMA, Volcan de Chiriqui (*Champion*).

This is a very distinct and pretty species; it is intermediate between those with the head sharply produced and those which have the vertex truncate, and not or scarcely produced before the eyes.