

(melal) = 3. **Ormenis erminia**, sp. n. (Tab. VII. figg. 20, 20 a.)

Alba vel pallide virescenti-alba, tegminibus ad costam et basim versus maculis vel lineis parvis fuscis instructis; capite perbrevis, oculis prominulis; pronoto vertice distincte longiori, leviter tricarinato; tegminibus apicem versus dilatatis, apice late et sat oblique truncato; corpore pedibusque albidis.

White or very pale greenish-white, with regular rows, of variable length, of small dark spots or dashes on the tegmina extending from the shoulders to behind the middle; head very short and without the eyes, which are prominent, distinctly narrower than the pronotum, which is longer than the vertex; pronotum tricarinate on the disc; tegmina broadest at the apex, the apex broadly and somewhat obliquely truncate; body and legs whitish.

Long. 15 millim.; lat. ad hum. 3 millim.

Hab. MEXICO, Teapa in Tabasco (*H. H. Smith*); PANAMA, Bugaba (*Champion*).

A specimen from Bugaba is figured.

4. **Ormenis leucophæa**.

Ormenis leucophæa, Stål, Stett. ent. Zeit. 1864, p. 55¹.

Hab. MEXICO (*Mus. Vind. Cæs.*, *Mus. Holm.*¹).

I cannot satisfactorily identify this species in our collection; we have two specimens from Atoyac, Vera Cruz, which may perhaps be discoloured examples of it.

5. **Ormenis infuscata**.

Ormenis infuscata, Stål, Stett. ent. Zeit. 1864, p. 55¹.

Hab. MEXICO (*coll. Signoret*¹, in *Mus. Vind. Cæs.*), Chilpancingo in Guerrero 4600 feet (*H. H. Smith*).

This is an obscure little species; I have referred one example in our collection to it, but somewhat doubtfully.

6. **Ormenis dolabrata**, sp. n. (Tab. VII. figg. 21, 21 a.)

Brunnea, variegata; capite testaceo, perbrevis, lato, cum oculis valde prominentibus pronoto distincte latiori, hoc et scutello testaceis, nigro sat late limbatis; tegminibus brunneis, plagâ magnâ testaceo-albidâ plus minusve irregulari ab humeris usque ad paullo ante apicem continuatâ, prope apicem marginem attingenti; alis plus minusve fumatis; corpore plus minusve ad latera virescente; pedibus testaceis.

Brown, variegated with whitish-testaceous; head and pronotum yellow-testaceous, the former very short and broad, eyes very large and prominent; pronotum dark at the sides, longer than the vertex; scutellum testaceous, with the sides broadly darker (the colour, however, of the front parts is very deceptive, and often appears to fade into a uniform brown after death); tegmina brown, with a broad whitish-testaceous irregular band reaching from the shoulders to a little before the apex, where it joins the margin; the claval suture and a broader or narrower line before the outer margin are also light-coloured, the colour being much more extended in some specimens than in others; wings smoky, darker towards the apex; abdomen greenish at the sides; legs testaceous.

Long. 9-10 millim.; lat. ad hum. 2 millim.

Hab. MEXICO, Cuernavaca and Miacatlan (*Bilimek*, in *Mus. Vind. Cæs.*), Chilpancingo in Guerrero 4600 feet (*H. H. Smith*).

A small series. At first I thought that these specimens might be referred to