

O. leucophæa, Stål, but they do not agree with the only typical example of that species which I have seen: this is in the Vienna Museum. I have, moreover, received examples of *O. dolabrata* among unnamed specimens from that collection. We figure an example from Chilpancingo.

7. *Ormenis squamulosa*, sp. n. (Tab. VII. figg. 22, 22 a.)

Fusca, plus minusve subcyaneo-pruinosa; capite testaceo, lato, perbrevis, oculis prominentibus; pronoto perbrevis, sed vertice longiori; scutello ad latera albido-squamoso; tegminibus marginibus supra et infra fere parallelis, apice subtruncato; pedibus testaceis

A rather small fuscous species, with a more or less distinct whitish-cyaneous deposit on the upper parts; head testaceous, vertex very short, eyes large and prominent; pronotum longer than the vertex; scutellum thickly squamose at the sides; tegmina with the upper and lower margins subparallel, and the apex somewhat obliquely truncate; body in fresh specimens more or less greenish; legs testaceous.

Long. $8\frac{1}{2}$ – $9\frac{1}{2}$ millim.; lat. ad hum. 2 millim.

Hab. MEXICO, Acapulco in Guerrero (*H. H. Smith*), Valladolid in Yucatan (*Gaumer*); GUATEMALA, San Juan in Vera Paz (*Champion*).

This species is very like *Pæciloptera glaucescens*, Walk., from San Domingo, but the head is differently shaped; it is also allied to *O. infuscata*, but is larger and has the head broader. A specimen from Acapulco is figured.

8. *Ormenis albescens*, sp. n. (Tab. VII. figg. 23, 23 a.)

Quoad formam staturamque præcedenti affinis, sed tegminibus totis albidis et capite angustiori distinguenda.

Very like the preceding in size and shape, but distinguished by having the tegmina entirely of a dirty-white colour and not squamulose, and by the narrower head; the front-parts are testaceous.

Long. 9 millim.; lat. ad hum. 2 millim.

Hab. MEXICO, Jalisco (*Schumann*).

9. *Ormenis griseoalba*, sp. n. (Tab. VII. figg. 24, 24 a.)

Griseo-albescens; capite perbrevis, lato, oculis prominulis; pronoto vertice distincte longiori; scutello longitudine fere latiori; tegminibus subfarinosis albescentibus, apicem versus plus minusve late fuscescentibus; corpore pedibusque albido-testaceis.

Of a greyish-white colour; head very short, broad, with the prominent eyes about as broad as the pronotum; pronotum longer than the vertex; scutellum almost broader than long; tegmina presenting a somewhat mealy appearance, more or less broadly infusate towards the apex; body and legs whitish-testaceous.

Long. 10 millim.; lat. ad hum. $2\frac{1}{2}$ –3 millim.

Hab. GUATEMALA, San Gerónimo (*Champion*); PANAMA (*Boucard*), David in Chiriqui (*Champion*).

This species is allied to *O. squamulosa*, from which it differs chiefly in its coloration and somewhat larger size. We figure an example from San Gerónimo.

NOTE.—There is a specimen of a rather fine species belonging to this group in our collection (12 millim. in length), from Cahabon, Vera Paz; it is, however, too imperfect to describe; the colour is dark grey, mottled with brown and whitish.