

2. *Tropidoscyta uniformis*, sp. n. (Tab. I. figg. 20, 20 *a*, 20 *b*.)

Præcedenti affinis, sed brevior, pronoto multo fortius punctato, carinisque lateralibus supra oculos distinctius elevatis; colore obscuriori, tegminibusque totis fuscis facillime distinguendus.

Very like the preceding, but smaller and shorter, black or almost black in colour, with the pronotum very coarsely punctured, and with the lateral carinæ above the eyes strongly marked; the tegmina are entirely fuscous, with the apex not hyaline, though slightly lighter than the base; the legs are fuscous.

Long. 4 millim.; lat. max. $2\frac{1}{2}$ millim.

Hab. PANAMA, Boquete, Bugaba, Volcan de Chiriqui, Caldera (*Champion*).

A specimen from Boquete is figured.

LEIOSCYTA, gen. nov.

Genus *Tropidoscytæ* affine, sed pronoto angustiori et longiori, carinisque lateralibus supra oculos deficientibus; tegminibus areis duabus discoidalibus, venisque minus elevatis instructis.

Allied to *Tropidoscyta*, but with the pronotum narrower and more elongate in proportion, and more parallel-sided if viewed from above; the upper surface, too, is smoother; the small carinæ over the eyes are wanting; the tegmina have only two discoidal areas, instead of three as in *Tropidoscyta*, and the veins are less raised and sometimes almost obsolete in the middle. The type of the genus is *Tropidoscyta pallidipennis*, Stål.

1. *Leioscyta pallidipennis*. (Tab. I. figg. 21, 21 *a*.)

Tropidoscyta pallidipennis, Stål, Kongl. Sv. Vet.-Ak. Handl. Band viii. 1, p. 46¹.

Hab. MEXICO¹ (*Mus. Holm.*; *Sallé*), Cuernavaca and Puebla (*Bilimek*, in *Mus. Vind. Cæs.*).

An example from the Stockholm Museum is figured.

2. *Leioscyta cornutula*. (Tab. I. figg. 22, 22 *a*.)

Tropidoscyta cornutula, Stål, Kongl. Sv. Vet.-Ak. Handl. Band viii. 1, p. 46¹.

Hab. MEXICO¹ (*Mus. Holm.*; *Sallé*).

The female only is known.

This species is closely allied to the preceding, but differs in having the front of the pronotum more porrect and extended into a very short blunt horn. It is quite possible that they may be sexes of one species; but I have only been enabled to examine one specimen of each.

3. *Leioscyta nitida*, sp. n. (Tab. I. figg. 23, 23 *a*, 23 *b*.)

Nigra vel fusco-nigra, nitida; pronoto fortiter punctato, carinâ centrali acute elevatâ plus minusve distincte albo-notatâ, et utrinque carinâ distinctâ sed minus elevatâ, instructo, lateribus supra oculos quoa dcarinas lævibus; tegminibus nigris ad medium opacis, apice abrupte et læte hyalino; pedibus nigris, genibus tarsisque dilutioribus vel totis fuscis.

A small, black, very shiny species, with the prothorax strongly and distinctly punctured, tricarinate, with the