

but it is not the "*semiclara*" of Signoret's Monograph, which is from Pulo Penang, as is evident from his description and figure (Ann. Soc. Ent. Fr. 1853, p. 666, t. 21. fig. 9).

14. ***Tettigonia melancholica***, sp. n. (Tab. XV. fig. 13.)

Præcedenti affinis, grisea vel brunneo-grisea; capite brevi, lato, oculis prominentibus; pronoto fasciâ latâ ad medium obscuriori; scutello griseo-testaceo; tegminibus isabellinis, fasciis tribus obscurioribus, apice hyalino; alis fumosis; abdomine supra rufo, corpore subtus testaceo; pedibus testaceis, plus minusve piceis.

In shape resembling the preceding species (*T. mosaica*), dun-coloured or griseous-brown; head short, broad, with prominent eyes; pronotum with a dark fascia situated just behind the middle; scutellum greyish-testaceous, with a broad fascia before and another behind the middle, and a narrower one before the subhyaline apex, darker; there are also two oblique darker lines, one on each side of the scutellum, and the veins are sometimes conspicuously darker than the ground-colour; wings smoky; abdomen red above; underside and legs testaceous, the latter more or less pitchy.

Long. 10-11 millim.; lat. ad hum. 3 millim.

Hab. PANAMA, Bugaba (*Champion*).

15. ***Tettigonia lineosa***, sp. n. (Tab. XV. fig. 14.)

Brevis oblonga, parallela, flava vel testacea; capite brevi, lato, rotundato, oculis prominentibus, nigro vittato; pronoto levissime transversim rugoso, marginibus et vittâ latâ transversâ ad medium nigris; scutello lineâ unâ transversâ nigrâ; tegminibus vittâ pone medium maculam albidam transversam includente, lineisque quibusdam obliquis a basi ad hanc extensis nigris, apice ipso fumoso; abdomine læte rufo; pedibus testaceis.

Short oblong, golden-yellow, yellow, or testaceous; head short and broad, rounded in front, with prominent eyes; metopidium black, with a larger or smaller central yellow patch, vertex black in front and behind, and with a transverse line across the middle; pronotum very finely transversely rugose, with the anterior margin narrowly and the posterior margin broadly black, and with a broad central median line; scutellum with a small central transverse line not reaching the margins; tegmina with a dark patch behind the middle enclosing towards its apex two whitish transverse sublunate spots, and with the margins, the suture, and several lines nearly or quite meeting this patch black; the line on the clavus is shorter; apex of the tegmina and the wings fuscous; abdomen bright red, with the central portion of the pygofer black; legs testaceous.

Long. 8 millim.; lat. ad hum. $2\frac{1}{2}$ millim.

Hab. PANAMA, Bugaba, Volcan de Chiriqui (*Champion*).

16. ***Tettigonia corixoides***, sp. n. (Tab. XV. fig. 15.)

Præcedenti similis, sed capite paullo longiori, magis producto; capite nigro, metopidio duabusque lineis notisque parvis pone ocellos flavis; pronoto nigro flavo-variegato, lævi; tegminibus antice hyalinis, venis fuscis, pone medium fuscatis hyalino variegatis; abdomine supra nigrescenti; corpore subtus cum pedibus dilute testaceis.

Like the preceding (*T. lineosa*) in general appearance, but with the head more produced and more angularly rounded; head black, with the metopidium and two transverse lines and two minute spots behind the ocelli sometimes confluent with the end of the hinder line yellow; pronotum black, with the central portion longitudinally and some patches towards the sides obscurely yellow; scutellum variegated; tegmina fuscous-hyaline in front, with the veins dark, and with the hinder part from a little beyond the