

middle fuscous, variegated with hyaline at the sides and before the apex; abdomen dark above; underside and legs light testaceous.

Long. 6-7 millim.; lat. ad hum. 2 millim.

Hab. PANAMA, Bugaba (*Champion*).

This species, as its name implies, bears a close resemblance at a little distance to a small species of *Corixa*.

17. *Tettigonia conspissata*, sp. n. (Tab. XV. fig. 16.)

Oblonga; capite pronotoque testaceis nigro figuratis, illo producto subtriangulâri late rotundato, oculis prominulis, hoc convexo basi haud sinuatâ; scutello figurato; tegminibus fuscis, lineis sex vel septem testaceis obliquis ad suturam vel ad tertiam partem apicalem extensis, hâc nigrescenti partim hyalinâ vittâ albidâ antice alterâque interruptâ postice instructâ; abdomine supra nigrescente; corpore subtus cum pedibus dilute testaceis, unguibus piceis.

Oblong, the head, pronotum, and scutellum testaceous, figured with black; head somewhat produced before the eyes, which are moderately prominent; pronotum convex, with the base not or hardly perceptibly sinuate; tegmina fuscous, with six or seven oblique testaceous lines extending from the shoulders to the apical third, which is dark, variegated with hyaline, and with two white or greenish-white bands which are made up of confluent spots and so not quite regular—one of these is situated at the front part of the dark portion and extends across the tegmina, the other (just before the apex) is interrupted at the suture; the hinder markings, however, appear to be somewhat variable; abdomen dark above; underside of body and legs light testaceous, the claws dark.

Long. 8-9 millim.; lat. ad hum. 2 millim.

Hab. GUATEMALA, Senahu and Chiacam in Vera Paz (*Champion*).

18. *Tettigonia multilineata*, sp. n. (Tab. XV. fig. 17.)

Sat lata, robusta, testacea; capite producto, triangulâri, oculis minus prominentibus, nigro figurato; pronoto brevi ut capite nigro figurato, leviter sculpturato; scutello nigro notato; tegminibus amplis, lineis sex vel septem obliquis utrinque instructis ad suturam vel ad apicem ipsum extensis; alis fumosis; abdomine subtus rufo, testaceo nigroque variegato; pedibus læte rufis, tarsis plus minusve fuscatis.

Rather a broad stout species, broadest at about the middle, testaceous, with the front parts figured with black; the tegmina with six or seven oblique lines on each parallel to the scutellum and reaching the suture on the apex, which is not hyaline; head produced, forming with the eyes, which are only slightly prominent, an equilateral triangle; pronotum rather short, slightly narrowed in front, gently sinuate at the base; tegmina smooth; underside of the abdomen red, variegated with black and testaceous, the front parts testaceous, variegated with black; legs bright red, with the tarsi more or less pitchy.

Long. 10-11 millim.; lat. ad hum. $3\frac{1}{2}$ millim.

Hab. MEXICO, Pinos Altos in Chihuahua (*Buchan-Hepburn*).

19. *Tettigonia lateralis*. (Tab. XV. fig. 18.)

Cercopis lateralis, Fabr. Ent. Syst., Suppl. p. 524¹.

Cercopis marginella, Fabr. Syst. Rhynch. p. 96².

Tettigonia costalis, Sign. Ann. Soc. Ent. Fr. 1854, p. 359, t. 12. fig. 8³.

Tettigonia lugens, Walk. List of Homopt. Ins. iii. p. 775⁴.

Tettigonia pyrrhotelus, Walk. loc. cit. p. 775⁵.

Hab. NORTH AMERICA, Canada², Nova Scotia⁴, Carolina¹, Florida⁵. — MEXICO
BIOL. CENTR.-AMER., Rhynch. Homop., Vol. II., August 1899.