

with the markings of the tegmina rather resembling those of *T. punctulata*, but it evidently belongs to this species.

We figure an example from Omilteme.

25. ***Tettigonia magica*.** (Tab. XV. figg. 24; 25, var.)

Tettigonia magica, Stål, Stett. ent. Zeit. xxv. p. 77¹.

Hab. MEXICO (*Mus. Holm.*¹ & *Mus. Vind. Cæs.*), Chilpancingo and Rincon in Guerrero (*H. H. Smith*), Cuernavaca in Morelos (*H. H. Smith, Bilimek*), Orizaba (*Bilimek, H. H. Smith, & F. D. Godman*).

This is a very easily recognized species, with the head rather produced and narrower than in several of the allied forms, and the markings of the head, thorax, and scutellum distinct and constant. I have found several specimens of it among unnamed material belonging to the Vienna Museum. As a rule, the tegmina have the ground-colour red, but a common variety has the whole ground-colour greenish-testaceous, without any red on the insect.

A specimen from Chilpancingo and a variety from Orizaba are figured.

26. ***Tettigonia consobrina***, sp. n. (Tab. XV. fig. 26.)

Fusco-rufa, maculis parvis dilutis regulariter aspersa; capite producto, haud triangulari, antice rotundato, oculis prominulis; pronoto sat longo, basi vix sinuatâ; scutello ad medium impresso; tegminibus apicem versus angustatis; abdomine supra fusco; corpore subtus rufescente; pedibus rufo-brunneis.

Dark fuscous-red, regularly sprinkled from the metopidium to the apex of the tegmina with small light spots, which are testaceous on the front parts, and greyish and more obscure on the tegmina; head produced, not triangular, rounded, eyes rather prominent; pronotum comparatively long; scutellum impressed in the middle; tegmina narrowed towards the apex; wings dark; abdomen fuscous above, underside reddish or reddish-yellow; legs reddish-brown, almost unicolorous, or with more or less of the tibiæ and tarsi infuscate.

Long. 7 millim.; lat. ad hum. 2 millim.

Hab. GUATEMALA, Cerro Zunil (*Champion*).

This species is allied to *T. punctulata*, but it is smaller, with the head longer and differently shaped, the pronotum longer in proportion, and the tegmina covered with small spots, whereas in *T. punctulata* they are more or less distinctly lined.

27. ***Tettigonia granulata*.** (Tab. XV. fig. 27.)

Tettigonia granulata, Walk. List of Homopt. Ins., Suppl. p. 195¹.

Hab. MEXICO (*Sallé, in Mus. Brit.*¹); GUATEMALA, San Gerónimo, Zapote, and Cerro Zunil (*Champion*).

We figure a specimen from San Gerónimo.