

47. **Tettigonia sirena.** (Tab. XVI. fig. 18.)

Tettigonia sirena, Stål, Stett. ent. Zeit. xxv. p. 76¹.

Hab. MEXICO (*Mus. Holm.*¹ & *Mus. Vind. Cæs.*).

This species is rather closely related to the preceding, and both are akin to *T. stål*; neither of them appears to be represented in our collection.

A specimen from the Vienna Museum is figured.

48. **Tettigonia albomaculata.** (Tab. XVI. fig. 19.)

Tettigonia albomaculata, Distant, Ent. Monthly Mag. xvi. p. 62¹.

Hab. COSTA RICA, Rio Sucio and Volcan de Irazu 6000 to 7000 feet (*Rogers*¹).

In this species the head and pronotum are black, and the tegmina are red, with a number of distinct white spots; the head is only a little produced before the eyes, and is broadly and regularly rounded in front.

A specimen from Rio Sucio is figured.

49. **Tettigonia tripunctata.**

Tettigonia tripunctata, Fitch, Fourth Ann. Report State Cab. Nat. Hist. p. 55 (1851)¹; Sign. Ann. Soc. Ent. Fr. 1854, p. 23, t. 2. fig. 12².

Tettigonia nigrifascia, Walk. List of Homopt. Ins. iii. p. 776³.

Tettigonia pallida, Walk. loc. cit. p. 776⁴.

Tettigonia albida, Walk. loc. cit. p. 777⁵.

Hab. NORTH AMERICA, New York¹.—MEXICO²⁻⁵ (*Mus. Vind. Cæs.*; *Boucard*, in *Mus. Holm.*), Omilteme in Guerrero, Cuernavaca in Morelos (*H. H. Smith*), Guanajuato (*E. Dugès*, in *Mus. Roy. Belg.*), Orizaba (*Bilimek*, *H. H. Smith*, & *F. D. Godman*); GUATEMALA, Quezaltenango 7800 feet, Dueñas (*Champion*).

This species varies considerably in colour and to a certain extent in the markings. It may be known from its allies by the very short head, which is scarcely produced before the eyes and is rounded in a very obtuse angle; the ground-colour is either reddish, greyish-brown, or yellowish-white. There is an insect in our collection from Mexico city which appears to be a variety of this species, but it has the head and pronotum covered with large black spots and markings; I cannot, however, separate it structurally.

50. **Tettigonia uniguttata.**

Tettigonia uniguttata, Walk. List of Hompt. Ins. iii. p. 778¹.

Hab. MEXICO¹.

This species differs from the preceding chiefly in having the head more conical, the vertex flatter, and the sculpture of the pronotum less evident; the head, pronotum,