

and tegmina are entirely testaceous, with the exception of a black spot on the front of the vertex: Signoret at first treated it as a variety of the preceding, but afterwards separated it. I am inclined to think that *T. uniguttata* is a colourless or immature variety of the preceding species.

51. ***Tettigonia præstantior***, sp. n. (Tab. XVI. fig. 20.)

Major, postice latior, capite producto, nigro, subexcavato, oculis sat elongatis; pronoto antice nigro postice læte flavescenti, leviter rugoso; scutello nigro; tegminibus lævibus, haud nitidis, basi et apice vittæque pone medium nigris, spatio intra vittam basalem et intermediam læte flavo, spatio post hanc flavo-rufo; corpore subtus plerumque testaceo; pedibus testaceis ad partem piceis.

A large, brightly-coloured species, with strongly contrasted black, yellow, and orange bands, the former having a cyaneous reflection in certain lights; head long, produced in an isosceles triangle, with the sides slightly rounded and bluntly angled at the apex, entirely black, except for a small portion of the metopidium above the rostrum—if viewed from the side, subexcavate, sharply pointed, and slightly deflexed; pronotum bright yellow, with the front part abruptly black, finely rugose; scutellum large, black; tegmina dull, smooth, with a black band at the base (forming with the scutellum, which it includes, a broad black band right across the body of the insect); a long yellow space behind this, and a broad black band just behind the middle, followed by a bright orange space before the apex, which is black; underside mostly testaceous, fuscous towards the apex of the abdomen; legs testaceous, more or less pitchy.

Long. 14–15 millim.; lat. ad hum. $3\frac{1}{2}$ millim.

Hab. PANAMA, Bugaba (*Champion*).

A considerable series. This is one of the most gaudily coloured and conspicuous insects of the whole group; it appears to be extremely local.

52. ***Tettigonia undecimmaculata***, sp. n. (Tab. XVI. fig. 21.)

Quoad staturam præcedenti affinis, sed capite breviori, oculis magis prominulis; testacea, capite nigro-notato, pronoto maculis tribus testaceis; tegminibus lævibus, sat nitidis, maculis octo conspicuis, duabus ad basim, quatuor ad medium, duabusque ante apicem instructis; corpore subtus pedibusque dilute testaceis.

Upperside fusco-testaceous, with black markings on the head and three light testaceous spots on the pronotum; head triangular, depressed, with the sides towards the front raised in a ridge on each side; tegmina smooth and rather shining, with two large round testaceous spots at the base, four in a line just behind the middle, the two inner ones being smaller and not rounded, and two large and conspicuous round spots at the apex; underside and legs light testaceous.

Long. 13 millim.; lat. ad hum. 3 millim.

Hab. PANAMA, Volcan de Chiriqui 3000 feet (*Champion*).

The specimen on which I have described this species appears to be somewhat immature, and the colour is probably darker as a rule, but the eight large and conspicuous spots on the tegmina render the insect so easily recognizable that I have thought it best not to pass it over.

53. ***Tettigonia feralis***, sp. n. (Tab. XVI. fig. 22, ♂.)

Fusco-nigra, capite magno, triangulariter producto; pronoto lævissime sculpturato, lateribus fere parallelis, basi vix sinuatâ; scutello magno; tegminibus lævibus, venis distinctis.