

58. *Tettigonia bilineata*, sp. n. (Tab. XVI. fig. 27.)

Aurea, lævis, nitida, capite fere semicirculari, maculâ nigrâ inter ocellos nigros rotundatâ vel cordiformi, oculis nigris vix prominulis; pronoto antice angustato, basi fere rectâ; scutello magno, concolori; tegminibus vittâ angustâ nigrâ curvatâ, scutellum attingente, et aliâ pone medium latiori instructis; abdomine flavo-rufo; pedibus dilute testaceis.

Golden-yellow, smooth, and shining, with the ocelli and a rather large round or heart-shaped spot between them, the eyes, which are scarcely prominent, and two bands on each tegmen, black; of the latter, the front ones reach the scutellum and extend in a curve to the costal margin, along which they are continued to near the middle of the tegmina; the hinder ones meet at the suture and form a straight, but more or less irregular band across the upper surface behind the middle; head rounded, nearly semicircular; pronotum narrowed in front; abdomen above orange-coloured; underside and legs light testaceous or rufo-testaceous.

Long. 10 millim.; lat. ad hum.  $2\frac{1}{2}$  millim.

*Hab.* MEXICO, Amula, Venta de Peregrino, Acaguizotla, and Rincon, all in Guerrero (*H. H. Smith*).

We figure a specimen from Amula.

59. *Tettigonia tæniata*, sp. n. (Tab. XVI. fig. 28.)

Aurea, nitida, lævis, vittis duabus latis cyaneo-nigris; capite lato, quadratim rotundato, vel fere semicirculari; ocellis et oculis nigris, his nonnihil prominentibus; corpore subtus pedibusque rufo-flavis; apice abdominis late, femoribus anticis superne cum tarsis, unguibusque intermediis et posticis nigris.

Of a bright golden colour, smooth, and shining, with two broad velvety cyaneous-black bands extending across the upper surface, the front one including the base of the pronotum, the scutellum, and the basal part of the tegmina, and the hinder one just before the apex, which is yellow; the apex of the abdomen is broadly black, and the upper surface of the front tibiæ, the front tarsi, and the intermediate and posterior claws are black.

Long. 11 millim.; lat. ad hum. 3 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

60. *Tettigonia distinguenda*, sp. n. (Tab. XVI. fig. 29.)

Præcedenti affinis, sed capite ad basim pronotoque ad apicem cyaneo-nigris, hoc parte basali flavescenti et tegminibus vittâ latâ ad basim scutellum includente nigrâ, alterâque apicem fere attingenti maculâ flavâ triangulari profunde indentatâ instructis, distinguenda.

Very like the preceding, of which it may possibly be a variety, but easily distinguished by the base of the head and front of the pronotum being black, as well as by the basal part of pronotum being yellow, and the apical band on the tegmina nearly reaching the apex, and being almost divided by a triangular yellow patch, the base of which rests on the margin.

Long. 11 millim.; lat. ad hum. 3 millim.

*Hab.* PANAMA (*Boucard*).

61. *Tettigonia ostrina*, sp. n. (Tab. XVI. fig. 30.)

Elongata, angusta, purpureo-rufa; capite pronotoque plus minusve dilutioribus, rufis vel rufo-flavis, illo sat producto obtuse rotundato, oculis prominulis, hoc convexo, lævi, antice angustato, basi leviter late sinuatâ; tegminibus lævibus sat nitidis, margine obscuriori, apice abrupte hyalino; abdomine supra rufo; corpore subtus pedibusque testaceis.

Elongate, narrow, of a purplish-red colour, with the head and pronotum more or less rufous or orange-coloured;

BIOL. CENTR.-AMER., Rhynch. Homop., Vol. II., February 1900.

\*33