

65. **Tettigonia trivirgata**, sp. n. (Tab. XVII. fig. 4.)

Elongata, parallela, purpureo-rufa; capite nigro antice miniato, triangulari, modice producto, oculis sat prominulis; fronte nigrâ; pronoto virgis tribus longitudinalibus læte miniatis; scutello magno; tegminibus maculis vel lineis quibusdam obscuris fusco-purpureis; corpore subtus pedibusque dilute testaceis.

Purple-red; head deep black, with the front part scarlet and the extreme apex with two black spots (which, if viewed from the front, form the upper portion of the black frons), short, triangular, moderately produced before the eyes, which are rather prominent; pronotum with three distinct, broad, scarlet, longitudinal bands; scutellum large; tegmina with the margins united a little before the apex, and a more or less elongate patch on each side of the suture obscurely fuscous-purple; underside and legs light testaceous.

Long. 8 millim.; lat. ad hum. vix 2 millim.

*Hab.* PANAMA, Bugaba and Volcan de Chiriqui (*Champion*).

This is a very distinct and pretty species, which may be known at once by the three scarlet bands on the pronotum.

We figure a specimen from Bugaba.

66. **Tettigonia purpurascens**, sp. n. (Tab. XVII. fig. 5.)

Elongata, parallela, purpureo-rufa, purpureo-fusco indistincte maculata; capite triangulari, ante oculos sat prominulos producto, late longitudinaliter sulcato; pronoto convexo, lateribus antice angustatis; tegminibus ad latera ante apicem maculis duabus parvis albidis, ad apicem ipsum hyalinis; corpore subtus pedibusque flavo-albidis, abdominis apice rufescenti.

Of a deep purple-red colour, with obscure lines of dark purple on the head and pronotum and spots of the same colour on the tegmina; head triangular, produced before the eyes; vertex broadly sulcate; metopidium light whitish-yellow; pronotum convex, with the sides gradually narrowed in front; scutellum large, somewhat raised behind; tegmina smooth, rather shiny, with two small white spots on each before the apex touching the margins, and the apex itself hyaline, fuscous at the sides; front portion of the underside and legs whitish-yellow.

Long. 8 millim.; lat. ad hum. 2 millim.

*Hab.* GUATEMALA, Cerro Zunil, Volcan de Atitlan (*Champion*).

The specimen figured is from the Volcan de Atitlan. Both localities are on the Pacific slope.

67. **Tettigonia costaricensis**. (Tab. XVII. fig. 6.)

*Tettigonia costaricensis*, Distant, Ent. Monthly Mag. xvi. p. 63<sup>1</sup>.

*Hab.* COSTA RICA, Rio Sucio and Volcan de Irazu 6000 to 7000 feet (*Rogers*<sup>1</sup>); PANAMA (*Boucard*).

This species may at once be known by its entirely black colour, with the exception of an orange band, which, including the pronotum (except just the front portion), encircles the whole body.

The specimen figured is from Panama.