

This insect appears to be locally abundant in various places in Central America.

Signoret² notes the fact that Latreille has given the length of the insect wrong in his description—24 millim. instead of 11 millim.

A specimen from David is figured.

71. *Tettigonia areolata*. (Tab. XVII. fig. 11.)

Tettigonia areolata, Sign. Ann. Soc. Ent. Fr. 1853, p. 355, t. 11. fig. 4¹.

Hab. MEXICO¹ (*Sallé*), Chilpancingo, Tierra Colorada, Dos Arroyos, and La Venta in Guerrero, Teapa in Tabasco (*H. H. Smith*), Atoyac in Vera Cruz (*Schumann*), San Lorenzo, Cordova (*M. Trujillo*), Temax in N. Yucatan (*Gaumer*); GUATEMALA, San Gerónimo (*Champion*); NICARAGUA, Chontales (*Janson*); COSTA RICA, Caché and Volcan de Irazu (*Rogers*), Candelaria Mts. (*Underwood*).

Signoret's figure of this species, as of several others more or less allied to it, is very poor and hardly recognizable; it is for this reason that we have thought it best to give fresh ones.

The specimen figured is from Tierra Colorada.

72. *Tettigonia collata*, sp. n. (Tab. XVII. fig. 12.)

Quoad staturam faciemque *T. areolatae* affinis; capite brevi, subtriangulari; metopidio rufo-flavo, brunneo-marginato; vertice antice rufo-flavo, postice nigro-brunneo vel brunneo; pronoto dilute flavo, marginibus anticis et posticis brunneis; tegminibus rufo-brunneis, maculis tribus utrinque dilute flavis vel albido-flavis; pedibus albido-flavescentibus.

Of about the size and shape of *T. areolata*; head short, subtriangular; metopidium orange, bordered with brown; vertex with the anterior half orange and the posterior half dark brown; pronotum light yellow, with the anterior and posterior margins dark brown; tegmina reddish-brown, with three light yellow patches on each, two large ones meeting behind the scutellum and not quite touching the margins, two just behind the middle forming a transverse band across the upper surface, and one smaller one on each side near the margin just before the apex, the apex not hyaline; legs very light yellow.

Long. 7 millim.; lat. ad hum. vix 2 millim.

Hab. PANAMA, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).

73. *Tettigonia laudata*. (Tab. XVII. fig. 13.)

Tettigonia laudata, Walk. Insecta Saundersiana, Homopt. p. 95¹.

Hab. MEXICO, Presidio de Mazatlan (*Forrer*), La Venta in Guerrero, Teapa in Tabasco (*H. H. Smith*), Valladolid in Yucatan (*Gaumer*); GUATEMALA, San Juan in Vera Paz (*Champion*); HONDURAS¹.

We figure a specimen from La Venta.

74. *Tettigonia sexguttata*. (Tab. XVII. fig. 14.)

Cicada sexguttata, Fabr. Syst. Rhynch. p. 75¹.

Tettigonia sexguttata, Sign. Ann. Soc. Ent. Fr. 1853, p. 357, t. 11. fig. 7²; Walk. List of Homopt. Ins., Suppl. p. 205³.