

base, one behind the middle, and a small one at the margin between these, the apex subhyaline; abdomen fuscous; sternum and legs light testaceous, apex of the posterior tibiae and the tarsi fuscous.

Long. 6 millim.; lat. ad hum.  $1\frac{1}{2}$  millim.

*Hab.* MEXICO, San Marcos (*Bilimek, in Mus. Vind. Cæs.*), Venta de Zopilote and Amula in Guerrero (*H. H. Smith*).

This species apparently belongs to the *T. venusta* group.

We figure a specimen from Venta de Zopilote.

### 85. *Tettigonia æqua*, sp. n. (Tab. XVII. fig. 23.)

Fusco-brunnea, antice albido et griseo variegata, tegminibus plus minusve obscure rufo vel griseo-rufo variegatis; capite producto obtuse rotundato, oculis vix prominulis; pronoto capite paullo longiori; tegminibus sat parallelis; abdomine supra fusco; corpore subtus et pedibus testaceis, plus minusve infuscatis.

Parallel, fuscous or fuscous-brown, with the head, pronotum, and scutellum variegated with lighter and darker shades of brown, as well as with greyish and testaceous colour; head produced, obtusely rounded in front; pronotum a little longer than the head; tegmina scarcely narrowed to the apex, with several irregular red or greyish-red spots on each, more or less obscure, apex subhyaline, with a little clear hyaline spot on each tegmen near the costal margin; abdomen fuscous above; underside and legs testaceous, more or less infuscate.

Long. 6-7 millim.; lat. ad hum.  $1\frac{1}{2}$  millim.

*Hab.* MEXICO (*Boucard, in Mus. Holm.*), Cuernavaca in Morelos (*H. H. Smith*); GUATEMALA, San Gerónimo (*Champion*).

This insect appears to be closely allied to *T. recta*, and at first I was inclined to regard it as a variety of that species; but the coloration is not the same and the head is a little differently shaped. The specimen from San Gerónimo may possibly belong to another species, but I do not feel justified in separating it.

An example from Cuernavaca is figured.

### 86. *Tettigonia albiceps*, sp. n. (Tab. XVII. fig. 24.)

Parva, fusca, fusco-rufo obscure notata; capite magno, producto, rotundato, albido, oculis prominulis; pronoto capite haud longiori, antice anguste albido; scutello ad medium late albido; tegminibus ad apicem obscure subhyalinis; abdomine supra late rufo; corpore subtus et pedibus albidis vel albido-testaceis.

A small, insignificant, fuscous species, with obscure red markings; head large, produced and rounded, about as long as the pronotum, whitish-testaceous, with two minute brownish spots at the base; forehead light brown, with two testaceous spots, one above the other, in the middle surrounded by dark colour; pronotum comparatively short, light in front, with two reddish spots on the anterior margin, joining the dark colour; tegmina obscurely suffused with red, especially at the claval suture; abdomen crimson above; underside and legs very light testaceous.

Long. 5 millim.; lat. ad hum. 1 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

This is an obscure little species, but it appears to be quite distinct, and may be easily known by its large, rounded, whitish head.