

91. **Tettigonia lutea.** (Tab. XVIII. fig. 1.)

Tettigonia lutea, Sign. Ann. Soc. Ent. Fr. 1855, p. 773¹.

Hab. MEXICO, Atoyac in Vera Cruz (*H. H. Smith*), Jalapa (*Bilimek*, in *Mus. Vind. Cæs.*); GUATEMALA (*Mus. Paris*¹), Cerro Zunil (*Champion*).

The hinder spot on the central line of the pronotum is sometimes wanting, and the tegmina, which are described by Signoret as "immaculées," occasionally have a small patch at the shoulder on a line within the outer margin fuscous.

The specimen figured is from Atoyac.

92. **Tettigonia hilaris.**

Tettigonia hilaris, Stål, Stett. ent. Zeit. xxv. p. 75 (1864)¹.

Hab. MEXICO (*coll. Signoret*¹; *Mus. Holm.*).

This appears to be merely a variety of the preceding (*T. lutea*) with two spots (instead of one) on the front of the vertex, and two (instead of one) towards the anterior margin of the pronotum; as, however, I have only seen one specimen I prefer not to unite them at present. The species is not represented in our collection.

93. **Tettigonia nigroguttata.**

Tettigonia nigro-guttata, Sign. Ann. Soc. Ent. Fr. 1855, p. 772, t. 23. fig. 8¹.

Hab. MEXICO (*Mus. Berol.*¹), San Marcos (*Bilimek*, in *Mus. Vind. Cæs.*).

This species is allied to the two preceding, but is larger and has a longer head. It is not represented in our collection.

94. **Tettigonia reservata**, sp. n. (Tab. XVIII. fig. 2.)

Subparallela, testaceo-viridescens; capite magno, producto, rotundato, dilute flavo, ocellis et maculâ ad medium verticis nigris; fronte, maculâ nigrâ ante verticem exceptâ, immaculatâ; pronoto virescente, antice testaceo, disco punctis duobus nigris transversim dispositis instructo; scutello magno, flavo-testaceo, concolori; tegminibus fere concoloribus; alis fumosis; abdomine supra nigro; corpore subtus pedibusque dilute flavo-testaceis.

Subparallel, greenish-testaceous; head large, produced, rounded, light yellow, with the ocelli and a spot on the middle of the vertex black; forehead unicolorous testaceous, with a black spot just before the vertex; eyes large and rather prominent; pronotum greenish, with the front broadly, and the rest of the margins narrowly, testaceous, the disc with two small black spots about the middle, arranged transversely; scutellum large, testaceous; tegmina dull, with a mealy appearance, the apex hyaline; wings dark; abdomen black above; underside and legs light yellow-testaceous, the claws fuscous.

Long. 6–6½ millim.; lat. 1½ millim.

Hab. MEXICO, Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*).

This species is allied to *T. lutea*, Sign., but differs in being shorter and dissimilarly coloured; it also has a black spot on the centre of the vertex, and two others (instead of one) on the disc of the pronotum.

The specimen figured is from Atoyac.