

95. ***Tettigonia 14-punctata*.**

Tettigonia 14-punctata, Sign. Ann. Soc. Ent. Fr. 1855, p. 782, t. 23. fig. 19¹.

Hab. MEXICO (*Mus. Berol.*¹).

A single specimen sent me as this species from the Vienna Museum has the head shorter, and broader in front, and the spots much larger, than shown in Signoret's figures. It appears to be merely a more spotted variety of *T. nigroguttata*.

96. ***Tettigonia lucasi*.** (Tab. XVIII. fig. 3.)

Tettigonia lucasi, Sign. Ann. Soc. Ent. Fr. 1855, p. 778, t. 23. fig. 16¹; Stål, Stett. ent. Zeit. xxv. pp. 76, 471².

Hab. MEXICO² (*Mus. Holm. & Mus. Vind. Cæs.*); GUATEMALA (*Mus. Paris*¹), San Gerónimo and Capetillo (*Champion*).

A specimen from San Gerónimo is figured.

97. ***Tettigonia edwardsi*.**

Tettigonia edwardsi, Sign. Ann. Soc. Ent. Fr. 1855, p. 780¹.

Hab. GUATEMALA (*Mus. Paris*¹).

I have not seen a specimen of this species, and it is not figured by Signoret.

98. ***Tettigonia dohrni*.**

Tettigonia dohrni, Sign. Ann. Soc. Ent. Fr. 1855, p. 793, t. 24. fig. 13¹.

Hab. MEXICO (*Mus. Berol.*¹).

I have not seen the type of this species, nor have I been able to identify it in our collection.

99. ***Tettigonia completa*, sp. n.** (Tab. XVIII. fig. 4.)

Flava, lineâ latâ centrali rufescente a fronte usque ad apicem tegminum extensâ, duabusque ad latera antice nigris instructa; capite magno, producto, oculis vix prominulis; pronoto convexo, nitido, haud capite longiori, basi leviter sinuatâ; tegminibus ad apicem angustatis; corpore subtus pedibusque flavis.

Yellow, with a broad central refescent line extending from the top of the forehead along the vertex (where it is dark in front), pronotum, scutellum, and suture to the apex of the tegmina; on each side also there is a corresponding line reaching from the metopidium along the margins of the pronotum (which are dark) and the sides of the tegmina; head as long as the pronotum, and a little narrower than its base, produced and rounded; pronotum convex and shining, very gently sinuate at the base; underside and legs yellow. Long. 7 millim.; lat. vix 2 millim.

Hab. MEXICO, Chilpancingo and Amula in Guerrero (*H. H. Smith*).

At first sight, this species resembles *T. lucasi*, but it differs entirely in the shape of the head, and the coloration is also distinct.