

Green, occasionally greenish-testaceous; head testaceous-yellow, somewhat produced and rounded, with two black lines nearly or quite meeting in front, two inside these meeting in an acute angle, and a central line at the base, often only indicated; all these lines extend on to the pronotum (which is testaceous in front, and has a small black line also at the margin), and the central three are continued on the testaceous scutellum; on each of the tegmina there are six or seven black lines, which are much more apparent on the lighter specimens; apex of the tegmina hyaline, with the extreme tip and a band before this fuscous; abdomen above black; underside and legs clear testaceous-yellow, the apex of the tibiae and claws being sometimes fuscous.

Long. 5-6 millim.; lat. $1\frac{1}{3}$ millim.

We figure a specimen from Bugaba.

129. ***Tettigonia sexlineata***. (Tab. XVIII. fig. 30.)

Tettigonia sexlineata, Sign. Ann. Soc. Ent. Fr. 1855, p. 792, t. 24. fig. 12¹.

Hab. MEXICO (*Bilimek*, in *Mus. Vind. Cæs.*), Orizaba (*Sallé*, *H. H. Smith*, & *F. D. Godman*), Cordova (*Sallé*), Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*); GUATEMALA (*Mus. Paris*¹), Chinautla 4100 feet (*Salvin*), Cerro Zunil (*Champion*); PANAMA, Bugaba (*Champion*).

The single specimen from Cerro Zunil has the black lines thicker and more pronounced than in our other examples.

We figure a specimen from Teapa.

130. ***Tettigonia sagata***.

Tettigonia sagata, Sign. Ann. Soc. Ent. Fr. 1854, p. 27, t. 2. fig. 15¹.

Hab. MEXICO (*coll. Germar & coll. Spinola*?¹).—BRAZIL, Rio Janeiro, Santa Teresa (*Mus. Roy. Belg.*).

I cannot identify this species in our collection, although there are several which are apparently very near it, to judge from Signoret's figure.

131. ***Tettigonia carissima***, sp. n. (Tab. XIX. fig. 1.)

Parva, virescenti-flava, nigro longitudinaliter vittata; capite sat lato, rotundato, oculis vix prominulis; fronte nigrâ; pronoto antice vix angustato; scutello magno; tegminibus ad apicem intus fumatis, iridescentibus; alis fuscis; abdomine fusco; pedibus albidis.

A very small species, greenish-yellow, with the metopidium black; on the upper surface two black stripes meet in the middle of the vertex, and are continued on each side nearly to the apex of the clavus, where they end in a point; between these markings there is a dark line extending along the centre of the pronotum and the scutellum; the sides of the tegmina are black, variegated with a few yellow stripes and patches, the yellow colour being continued to the apex, the inside apical portion being smoky and iridescent; wings and abdomen infusate; legs slender, whitish.

Long. $4\frac{1}{2}$ -5 millim.; lat. ad hum. 1 millim.

Hab. PANAMA, Bugaba (*Champion*).

This very pretty little species appears to be considerably smaller than *T. sagata*; the head, too, is more rounded, and has the two longitudinal black stripes meeting in