

and succeeding species; abdomen crimson above, testaceous beneath; sternum in part black; legs testaceous, femora broadly black towards the base.

Long. 5 millim.; lat. $1\frac{1}{2}$ millim.

Hab. MEXICO, Chilpancingo in Guerrero (*H. H. Smith*).

150. *Tettigonia robustula*, sp. n. (Tab. XIX. fig. 21.)

Parva, robusta; capite, pronoto, scutelloque testaceis; fronte nigro-lineatâ, maculâ magnâ nigrâ ante verticem, vertice producto, triangulari, oculis nigris prominulis; tegminibus sat latis, brunneo-testaceis unicoloribus; alis latis fumosis, apicibus truncatis; abdomine nigro; sterno testaceo, nigro-variegato; pedibus dilute testaceis, apice tibiæ et tarsis plus minusve infuscatis.

A small, obscure, stoutly-built species, with the front-parts testaceous, more or less suffused with obscure darker markings, especially on the pronotum; forehead lined with black and with the usual spot before the vertex well pronounced; vertex triangular, considerably produced and plainly angled in front; eyes black, rather prominent; pronotum shorter than the head, convex, narrowed in front, scarcely, if at all, sinuate at the base; tegmina comparatively broad, brownish-testaceous, narrowly hyaline at the apex; wings smoky-brown, truncate at the apex; abdomen black, more or less ringed with yellow, especially below; sternum testaceous, variegated with black; legs testaceous, with a dark spot at the base of the femora and with the apex of the tibiæ and the tarsi more or less impunctate.

Long. 5 millim.; lat. $1\frac{1}{2}$ millim.

Hab. MEXICO, Sierra de las Aguas Escondidas and Omilteme in Guerrero, 8000 to 9000 feet (*H. H. Smith*).

151. *Tettigonia satelles*, sp. n. (Tab. XIX. fig. 22.)

Parva, nigra, capite obtuse triangulari, oculis leviter prominulis; pronoto antice angustato; capite vix longiori; scutello magno; tegminibus maculis quatuor, duabus ad marginem costalem, maculâ hyalinâ divisis, et duabus ad marginem interiorem instructis; alis fumosis; abdomine supra nigro; corpore subtus et pedibus albidis.

A small, black, rather shining species; head obtusely triangular, with the eyes slightly prominent; forehead light testaceous, without any spot; pronotum almost as long as the head, very finely rugose transversely, with the base nearly straight; scutellum rather large; each tegmen with four white waxy-looking spots—two on the costal margin and two on the interior margin, the front ones of the latter being semicircular and so forming a round comma-shaped spot when the tegmina are closed; between the two costal spots is a hyaline patch; abdomen black above; underside and legs pale whitish-testaceous, clavus and sometimes part of the tarsi a little infuscate.

Long. 4 millim.; lat. 1 millim.

Hab. MEXICO, Teapa in Tabasco (*H. H. Smith*).

This species is allied to *T. quadriplagiata*, Walk., from the "West Coast of America," but that insect has a red abdomen and different markings. Teapa, it may be noted, is on the Atlantic slope.

152. *Tettigonia mollicula*, sp. n. (Tab. XIX. fig. 23.)

Parva, albida, fusco-lineata; capite sat magno, producto, rotundato, oculis haud prominulis; fronte dense nigro-lineatâ, maculâ nigrâ ante verticem, vertice lineâ latâ albidâ longitudinali fusco limitatâ; pronoto subparallelo, vittis duabus latis longitudinalibus rufo-brunneis ad medium, et duabus angustis fuscis ad latera instructo; scutello lineâ centrali fuscâ; tegminibus subvitreis, distincte fusco lineatis; alis vitreis; abdomine supra nigro; corpore subtus et pedibus albidis.